

# Robust Sensitivity Analysis for Stochastic Systems

Henry Lam\*  
*Boston University*

## Abstract

Sensitivity analysis for stochastic systems is typically carried out via derivative estimation, which critically requires parametric model assumptions. In many situations, however, we want to evaluate model misspecification effect beyond certain parametric family of models, or in some cases, there plainly are no parametric models to begin with. Motivated by such deficiency, we propose a sensitivity analysis framework that is parameter-free, by using the Kullback-Leibler divergence as a measure of model discrepancy, and obtain well-defined derivative estimators. These estimators are robust in that they automatically choose the worst (and best)-case directions to move along in the (typically infinite-dimensional) model space. They require little knowledge to implement; the distributions of the underlying random variables can be known up to, for example, black-box simulation. Methodologically, we identify these worst-case directions of the model as changes of measure that are the fixed points of a class of functional contraction maps. These fixed points can be asymptotically expanded, resulting in derivative estimators that are expressed in closed-form formula in terms of the moments of certain “symmetrizations” of the system, and hence are readily computable.

## 1 Introduction

The evaluation of performance measures for stochastic systems, be it using simulation or analytical techniques, often relies heavily on the underlying model assumptions. Typically, the computational goal is the expectation of a certain functional  $h(\cdot)$  that depends on the underlying random components, say,  $\mathbf{X} = (X_1, X_2, \dots)$ , i.e.  $E[h(\mathbf{X})]$ . The model assumptions on these random components intricately affect the quantity to be estimated. For example, in queueing networks, the expected queue length and workload depend on the distributional assumptions imposed on the sequence of interarrival and service times. In finance, the computation of option prices requires the transition probability distributions of the asset price movement and risk factors. Computations that are based on wrong assumptions on the underlying components can affect the output in various magnitudes.

To address this issue, a vast literature has been written to assess the effect of model misspecification. One major assessment tool is sensitivity analysis, or more specifically derivative estimation, by calculating the effect on the performance measure due to an infinitesimal change of the specified parameter in the model assumption. The computation of such derivative estimators can be done via several techniques. The most straightforward method is to use finite difference and invoke the first principle of differentiation [15, 53]. When the dependency of the parameter is in the functional  $h$ , under some regularity conditions, differentiation and expectation operators can be interchanged to obtain derivative estimators directly in expectation form; this is known as pathwise differentiation or infinitesimal perturbation analysis (see, for instance, [28, 18, 19]). On the other hand,

---

\*Department of Mathematics and Statistics, Boston University. Email: [kh1am@bu.edu](mailto:kh1am@bu.edu).

when the parameter is involved in the density function, the likelihood ratio method (or the score function method) can be employed via differentiation of the density, resulting in an estimator in the form of an expectation that is weighted by the so-called score function (see [23, 44]). Other techniques that seek to improve and generalize the above standard methods include conditioning and smoothed perturbation analysis [17, 24], kernel estimator [38], the push-out method [46, 47], and other modifications or combinations of the above [32, 52, 34, 45, 29]. The surveys [33], [16], §VII in [3] and §7 in [20] provide general overview on these different methods.

All these derivatives estimators, both conceptually and practically, require to certain extent a parametric model to establish. Of course, when the model is known to belong to a particular family, they can be effective in the assessment. However, more often than not, we want to assess the effect on our estimate due to model discrepancy that potentially goes beyond a change in a specified parameter. For example, a communication network administrator wants to forecast the chance of capacity overflow, but may not know exactly if Markovian assumption holds for the arrival and service time distributions. A portfolio manager who wants to predict future return may have doubt about the lognormal assumptions made on the individual asset price movements. Assessing the effect of model deviations of this sort clearly goes beyond the classical parametric derivative framework. More importantly, in many instances in operations management, there is no natural guidance in deciding a “true” parametric model; rather, nonparametric methods based on past data are adopted. In these situations there is not even a parametric family to begin with, not to mention the issue of assessing the effect of model uncertainty via classical sensitivity analysis.

With such motivation, we develop a notion of *nonparametric derivatives* that can assess, in a well-defined sense, how the performance measure changes under an infinitesimal change in the general model space. These nonparametric derivatives possess the following features:

1. **Parameter-free/data-driven.** Minimal information about the underlying model is required. For example, the distribution of the underlying random components can be known up to black-box simulation. They can also be known in the form of smoothed empirical distributions based on collected data, or other primitive forms. In particular, no parameter is needed for any of our analysis.
2. **Robust/adversary.** These estimators capture model discrepancies along the worst and best-case directions in the space of admissible models (which is typically infinite-dimensional). As a result, they dominate all classical parametric derivatives (in a sense we will discuss).
3. **Computationally efficient.** To compute these estimators, no additional knowledge is required other than the numerical generation of the random components and the functional of interest. In fact, the form of these estimators depends only on the moments of certain “symmetrizations” of the stochastic system, via a simple sequential conditioning scheme, and hence they are readily computable.

Let us briefly explain our methodology for developing these nonparametric derivatives; the main idea and formulation will be presented in more details in Section 2. First, to properly define derivatives one needs a notion of distance. Under parametric assumptions, Euclidean distance is naturally employed as a measurement of distance. In the nonparametric situation, one can use the notion of statistical distance [51] defined between two probability distributions, such as Kullback-Leibler (KL) divergence [31], to measure the closeness of models. However, without parametric assumptions, the space of admissible models is generally infinite-dimensional; consequently, blindly shrinking the statistical distance to zero and applying the first principle of differentiation can lead

to different rate-of-change values that result from different directions of movement. The key, then, is to use a *robust* approach and find the natural steepest descent direction among all admissible directions. Doing so requires solving appropriately posted optimization problems, which give the best and worst-case directions of movement of model discrepancy, before shrinking the distance to zero.

These optimization problems, which we will describe in Section 2, have strong connection to the robust control and optimization literature. Roughly speaking, the goal of this literature is to design the notion of optimum and to find optimal decisions in contexts where full probabilistic characterization is unavailable or intractable. One common approach is to consider max min objective, where the minimum is taken over a class of models that is believed to contain the truth, often called the uncertainty set [35]. The choice of the class defining the uncertainty set can vary, depending on both belief and computational consideration. In particular, a class that has been extensively studied is deterministic robust optimization, where the model is taken as degenerate and the uncertainty set is a deterministic region (see for instance [6, 7, 5, 4]). When the model in consideration is not deterministic, the uncertainty set is usually taken as a region defined via statistical distance such as KL divergence. This approach has also been studied widely, particularly for control problems in engineering [43], economics and finance [26, 27, 21, 37], queueing [30], and dynamic pricing [36]. As a tool for model sensitivity assessment, our approach naturally takes the latter type, i.e. the discrepancy in model is probabilistic. Rather than searching for an optimal control policy, however, our goal is to assess the effect of model discrepancy. In this regard, [22] has considered a similar formulation and coined the technique *robust Monte Carlo* as the solution to a class of convex programs that arise in finance (the reason that *Monte Carlo* appears is because of a naturally defined importance sampling scheme for the dual problem; see (26) below). This solution, as we will discuss, is a consequence of single-stage robust control and serves as an important building block for our development.

Despite such a rich literature on robust optimization, the derivation of our results calls for new methodological development, for the following reason: the behavior of derivative estimators relies crucially on the *local*, not global, properties of the associated optimization problems. In fact, with the assumptions on  $h$  that we shall discuss, our posted optimization problems are in general non-convex and globally intractable. This issue is especially prominent in the case of multi-stage problems, when a sequence of random variables, all generated according to the same model, shows up in the evaluation of the performance measure; such scenario appears often in real applications (think of, for example, the sequence of interarrival times and service times in a queue generated in an i.i.d. fashion). Therefore, instead of directly solving these optimization problems, we leverage the stochastic interpretations on the underlying system to carry out local analysis. In particular, to characterize the steepest descent directions, we develop a new representation of the associated locally optimal changes of measure as the fixed points to a class of functional contraction maps. These fixed points in turn possess tractable asymptotic behavior as the involved statistical distances vanish, and support high order Taylor series type expansions. We shall highlight that, in addition to obtaining asymptotic expansions, the contraction characterization of the optimal changes of measure is of interest itself and gives a precise description on the optimal movement under the involved structural landscape. The study of these contraction operators and their fixed points will be central in our line of mathematical development and drive the key towards the intriguing “symmetrization” form of our derivative estimators.

Lastly, our development in this paper relates to two lines of study in the statistics literature, one on the form of our estimators and the other on our formulation; we believe that a deeper

understanding in both connections, with the goal of integration with real data, is important and shall comprise future research. Here we briefly discuss the relations and the differences between our development and these lines of study. The first one is the asymptotic distribution theory of von Mises differentiable statistical functions (§6 in [48]). This theory studies the rate of convergence and the normality properties of statistics that can be expressed as functionals of empirical distributions. There, the main development involves expansions using Frechet derivatives that reduce the problem into analyzing some  $U$ -statistics of the individual samples, whose asymptotic moments turn out to bear a “symmetrization” form similar to this paper. In the related literature of robust statistics, such symmetrization form can be interpreted as the so-called “influence curve” that measures the effect to a statistic due to contamination at each arbitrary point in the space [25]. Methodologically, however, the derivation in von Mises’ theory crucially utilizes Levy’s continuity theorem and spectral analysis, in contrast to the optimization and change-of-measure approach proposed in this paper. Such methodological difference arises as von Mises’ theory concentrates on finding central limit asymptotics for statistics of interest as data size grows, whereas our focus is on quantifying the effect of model discrepancy using a robust formulation without a priori data structure. We shall leave the full exploration of this connection in future research, which we believe can be fruitful in gaining further understanding on data-related robustness properties of some important performance measures in the operations research literature.

The second related line of study is the empirical likelihood (EL) method [42]. This is the revisit of classical inference problems, such as hypothesis testing, that uses empirical distributions instead of parametric models. Because the empirical distribution is the maximum likelihood estimator in the space of distributions, an analog of Wilks’ Theorem ([12] p. 92) establishes that the associated logarithmic likelihood ratio between the empirical distribution and the true hypothesis (known as the profile likelihood) is asymptotically chi-square. Consequently, the computation of confidence region for a certain model parameter (such as moment) can be formulated as a constrained optimization problem with the associated critical values. The constraint of this optimization problem, which involves the so-called empirical likelihood, is analogous to the uncertainty set in the robust optimization formulation, and finding the confidence region is analogous to finding the range of performance outputs between the best and worst-case directions of model discrepancy in our setting. Although the EL method involves quite different asymptotic calculations due to its focus on inference, it can potentially be combined with our nonparametric derivative machinery to construct asymptotically valid confidence intervals for simulation estimators that are subject to input model uncertainty; we shall provide some discussion of this connection in Section 8.

This paper is organized as follows. First, we explain our main idea and formulation in Section 2. We then state our main results in Section 3. Methodological development will be presented in Sections 4 and 5, followed by a discussion on the connection to parametric models in Section 6. Section 7 will then demonstrate our methods in some numerical experiments, and Section 8 will be devoted to a discussion of future work. We defer some technical proofs to the appendix.

## 2 Main Idea and Formulation

To explain our framework, let us call a real-valued function  $h(\cdot)$  the *cost function* for convenience, and we are interested in the *performance measure*  $E[h(\mathbf{X})]$ , where  $\mathbf{X} = (X_1, X_2, \dots)$  is a (potentially infinite) sequence of i.i.d. random components each lying on the domain  $\mathcal{X}$ . The cost function  $h$  and the domain  $\mathcal{X}$  can be quite general. For example,  $h(\mathbf{X})$  can be the waiting time of the 100-th customer in a queueing system, where  $\mathbf{X}$  is the sequence of interarrival and service times.

Our nonparametric sensitivity analysis starts from a *benchmark* model that is believed to describe the random components  $X_1, X_2, \dots$ , and is used in the primary computation. Denote  $P_0$  and  $E_0$  as the probability distribution and the expectation under this benchmark model. We call  $E_0[h(\mathbf{X})]$  the benchmark performance measure. The goal is to assess the effect on  $E_0[h(\mathbf{X})]$  when there is an infinitesimal change in the model, i.e. the distribution  $P_0$ .

As discussed in the introduction, to properly define derivatives in nonparametric settings, we use the notion of statistical distance defined on probability distributions as a measurement of model discrepancy. Throughout this paper we shall use KL divergence, defined between two distributions  $P_1$  and  $P_2$ , as

$$D(P_1 \| P_2) := \int \log \frac{dP_1}{dP_2} dP_1 = E_2 \left[ \frac{dP_1}{dP_2} \log \frac{dP_1}{dP_2} \right]$$

where  $dP_1/dP_2$  is the likelihood ratio, equal to the Radon-Nikodym derivative of  $P_1$  with respect to  $P_2$  (assuming absolute continuity), and  $E_2[\cdot]$  is the expectation under  $P_2$ . The choice of KL divergence has several analytical advantages over other statistical distances and streamlines our development, as will be seen in the next section. Although KL divergence is, strictly speaking, not a metric, it serves as a measurement of distance in that two distributions possess similar statistical properties when they have a small KL divergence [51]. Supposing there is a slightly perturbed version of the benchmark model  $P_0$ , say  $P_f$ , so that  $D(P_f \| P_0) \approx 0$ , indeed one would expect that  $E_f[h(\mathbf{X})] \approx E_0[h(\mathbf{X})]$ , which provides a basis for a rate-of-change analysis.

The main issue with plainly using the first principle of differentiation, when shrinking  $D(P_f \| P_0)$  to 0, is the dimensionality: the space is now the set of probability distributions that are absolutely continuous with respect to  $P_0$ , which is typically infinite-dimensional. As a result, moving along different directions in this space can lead to different rates of change in the performance measure. To tackle this issue, as discussed in the introduction, we shall adopt a robust approach by considering the most extreme directions in taking the derivative, namely the ones that give the highest and the lowest rates of change in the performance measure. More concretely, before shrinking the KL divergence  $D(P_f \| P_0)$  to 0, consider the optimization problems

$$\begin{aligned} \max \quad & E_f[h(\mathbf{X})] \\ \text{subject to} \quad & D(P_f \| P_0) \leq \eta \\ & P_f \in \mathcal{P}_0 \end{aligned} \tag{1}$$

and

$$\begin{aligned} \min \quad & E_f[h(\mathbf{X})] \\ \text{subject to} \quad & D(P_f \| P_0) \leq \eta \\ & P_f \in \mathcal{P}_0 \end{aligned} \tag{2}$$

where  $\mathcal{P}_0$  is the set of all probability distributions that are absolutely continuous with respect to  $P_0$ , which comprises our *admissible* model space. The problems (1) and (2) calculate the maximal and minimal values of the performance measure when the perturbed distribution  $P_f$  is within  $\eta$ -neighborhood of  $P_0$ .

The key message of this paper is that when letting  $\eta$  go to 0, under mild assumptions on  $h(\mathbf{X})$ , the optimal values of (1) and (2) can each be expressed as

$$E_{f*}[h(\mathbf{X})] = E_0[h(\mathbf{X})] + \zeta_1(P_0, h, \mathbf{X})\sqrt{\eta} + \zeta_2(P_0, h, \mathbf{X})\eta + \dots \tag{3}$$

where  $\zeta_1, \zeta_2, \dots$  is a sequence of coefficients that can be written in closed-form formula in terms of  $h$  and  $\mathbf{X}$  under the benchmark distribution  $P_0$ . The coefficient  $\zeta_1$  can be interpreted as the first order

nonparametric derivative,  $2\zeta_2$  as the second order, and so forth, as a mimic of Taylor series. The rest of this paper focuses on finding these coefficients, presented in a roughly increasing order of sophistication regarding system complexity. Also, let us concentrate on the maximization problem (1), since the minimization counterpart (2) is merely a replacement of  $h$  by  $-h$ ; we shall discuss this implication in Section 6.1.

### 3 Main Results

We state our results on three levels, namely when the cost function  $h$  depends on a single variable, a sequence of i.i.d. variables on a fixed time horizon, and a sequence of i.i.d. variables on a random time horizon.

#### 3.1 Preliminary Results and Insights

Suppose that the cost function  $h$  depends only on a single variable  $X \in \mathcal{X}$  in the formulation (1), i.e. the maximization problem (1) is

$$\begin{aligned} \max \quad & E_f[h(X)] \\ \text{subject to} \quad & D(P_f \| P_0) \leq \eta \\ & P_f \in \mathcal{P}_0. \end{aligned} \tag{4}$$

We impose the following assumptions:

**Assumption 1.** *The variable  $h(X)$  has exponential moment in a neighborhood of 0 under  $P_0$ , i.e.  $E_0[e^{\theta h(X)}] < \infty$  for  $\theta \in (-r, r)$  for some  $r > 0$ .*

**Assumption 2.** *The variable  $h(X)$  is non-degenerate, i.e. non-constant, under the benchmark distribution  $P_0$ .*

The first assumption is on the light-tailedness of  $h(X)$ . This is in particular satisfied by any bounded  $h(X)$ , which handles all probability estimation problems. The second assumption is to exclude the uninteresting degeneracy case. More importantly, it also ensures that the benchmark distribution  $P_0$  is not an optimal model, since non-degeneracy always leads to an opportunity of upgrading the performance output by rebalancing the measure. This in turn guarantees the validity of an ascent direction. In this single-variable scenario, we can get a very precise understanding of the effect of model misspecification when  $\eta$  is small:

**Theorem 1.** *Suppose Assumptions 1 and 2 hold. Denote  $\psi(\beta) = \log E_0[e^{\beta h(X)}]$  as the logarithmic moment generating function of  $h(X)$ . When  $\eta > 0$  is within a sufficiently small neighborhood of 0, the optimal value of (4) is given by*

$$E_{f^*}[h(X)] = \psi'(\beta^*) \tag{5}$$

where  $\beta^*$  is the unique positive solution to the equation  $\beta\psi'(\beta) - \psi(\beta) = \eta$ . In particular, we have

$$E_{f^*}[h(X)] = E_0[h(X)] + \sqrt{2\text{Var}_0(h(X))}\eta^{1/2} + \frac{1}{3} \frac{\kappa_3(h(X))}{\text{Var}_0(h(X))}\eta + O(\eta^{3/2}) \tag{6}$$

where  $\text{Var}_0(h(X))$  is the variance of  $h(X)$ , and  $\kappa_3(h(X))$  is the third order cumulant of  $h(X)$  under  $P_0$ .

Below are some observations concerning Theorem 1 that can be shown in significant generality:

1. *First order derivative interpretation.* First, from (6) it appears that the square root of KL divergence is a correct scaling of the first order model misspecification effect. The first order derivative of  $E_{f^*}[h(X)]$  with respect to  $\sqrt{\eta}$  is

$$\lim_{\eta \rightarrow 0} \frac{E_{f^*}[h(X)] - E_0[h(X)]}{\sqrt{\eta}} = \sqrt{2\text{Var}_0(h(X))}$$

which can be interpreted as the maximal *first order nonparametric derivative* with respect to KL divergence.

2. *Second order derivative interpretation.* Similar to the first order derivative, the quantity  $(2/3)(\kappa_3(h(X))/\text{Var}_0(h(X)))$  can be interpreted as the *second order nonparametric derivative*. It can be written more explicitly in terms of higher order moments as

$$\frac{1}{\text{Var}_0(h(X))} \left[ \frac{2}{3}E_0[h(X)^3] - 2E_0[h(X)]E_0[h(X)^2] + \frac{4}{3}(E_0[h(X)])^3 \right]$$

3. *Form of the derivatives.* It turns out that the form  $\sqrt{2\text{Var}_0(h(X))}$  is a general pattern: even in more complex systems, the first order derivative is always of the form  $\sqrt{2\text{Var}_0(g(X))}$  where  $h$  is replaced by a function  $g$  that captures some kind of “symmetrization” of  $h$ . This symmetrization typically involves a sum of conditional expectations. In the case of random time horizon problems, the sum can be infinite (yet still computable). The second order derivative can involve an extra term that captures a second order “symmetrization” of  $h$ . These generalizations will be discussed in Sections 3.2 and 3.3.
4. *Translation invariance.* Both the first and the second order nonparametric derivatives are translation invariant in  $h(X)$ , i.e. they retain the same values when  $h(X)$  is replaced by  $h(X) + c$  for some constant  $c$ . This is apparent from the formula in (6) since cumulants are invariant to constant shifting. It is also consistent with the intuitive notion that adding a constant to  $h(X)$  should not change its sensitivity analysis. This observation has implication in the development of the corresponding quantities in random time horizon problems in Section 5.5.
5. *Sign of the derivative.* The first order nonparametric derivative  $\sqrt{2\text{Var}_0(h(X))}$  is always positive. This is because of the maximization formulation. The derivative is taken along the direction, among all the eligible changes of measure, that gives the largest increase in the expectation of the cost function. In general, when the formulation is a minimization, the sign of the first order derivative is negative while keeping the same magnitude, and the second order derivative is the same as the maximization counterpart. Under Assumptions 1 and 2, the only change in (5) for the minimization formulation is that  $\beta^*$  should be the unique negative solution of the same equation.
6. *Parametric dominance.* Because of the optimization formulations (1) and (2), the magnitudes of our nonparametric derivatives always dominate any eligible classical parametric derivative, under a rescaling from Euclidean distance to KL divergence. This will be discussed in more details in Section 6.2. Example 1 below also illustrates this point.

Before we move on, let us consider a toy example for illustration:

**Example 1** (Gaussian mean). Consider the quantity  $EX$ , where  $X$  is a mean-zero Gaussian random variable with variance  $\sigma^2$ . Then obviously  $E_0X = 0$ . We will compare a classical parametric derivative with our nonparametric approach. First, suppose the mean of  $X$  is misspecified, but it is still Gaussian with variance  $\sigma^2$ , so that the mean is  $\mu \neq 0$  instead of 0, i.e.  $E_fX = \mu$ . This model misspecification effect can be summarized by establishing that the first order parametric derivative of  $EX$  with respect to  $\mu$  is 1.

To quantify this movement of  $\mu$  in terms of KL divergence, let us consider  $P_f$  as  $N(\mu, \sigma^2)$  in the Gaussian family. It is an easy exercise to derive the relation  $D(P_f||P_0) = \mu^2/(2\sigma^2)$ :

$$D(P_f||P_0) = E_0[L \log L] = E_0 \left[ \exp \left\{ \frac{1}{\sigma^2} \left( Z\mu - \frac{\mu^2}{2} \right) \right\} \frac{1}{\sigma^2} \left( Z\mu - \frac{\mu^2}{2} \right) \right] = \frac{\mu^2}{2\sigma^2}$$

where  $L = \exp \left\{ \frac{1}{\sigma^2} \left( Z\mu - \frac{\mu^2}{2} \right) \right\}$  is the likelihood ratio between  $P_f$  and  $P_0$ , and  $Z$  denotes a  $N(0, \sigma^2)$  random variable. Hence a movement of one unit of  $\mu$  triggers a movement of  $\mu^2/(2\sigma^2)$  units in KL divergence. In terms of the derivative with respect to the square root of KL divergence, the first order sensitivity of  $E_fX$  is then  $1 \times \sqrt{2}\sigma = \sqrt{2}\sigma$  along the direction of movement in mean within the Gaussian family with variance  $\sigma^2$ .

Going to our formulation, Theorem 1 states that the maximal increase in the value of  $EX$ , over all admissible distributional movement, is

$$E_{f^*}X - E_0X = \sqrt{2\text{Var}_0(X)}\eta^{1/2} + O(\eta) = \sqrt{2}\sigma\eta^{1/2} + O(\eta).$$

In particular, according to discussion point 6 above (which will be justified rigorously in Proposition 4), this should provide an upper bound for the sensitivity when confined to Gaussian model. Indeed, the first order nonparametric derivative is  $\sqrt{2\text{Var}_0(X)} = \sqrt{2}\sigma$ , which matches exactly with the parametric derivative along the movement in mean within the Gaussian family.

In the coming subsections we shall generalize the above observations to more complex systems where sequences of random components are involved.

### 3.2 Finite Horizon Problems and Symmetrization Estimators

In this section, we consider cost functions that depend on not only one but a sequence of random variables. We focus on the case when the sequence is i.i.d. on a fixed time horizon. Namely, we consider  $E[h(\mathbf{X}_T)]$ , where  $\mathbf{X}_T = (X_1, \dots, X_T) \in \mathcal{X}^T$ , each  $X_t$  being i.i.d., for the fixed time horizon  $T > 1$ . For convenience we also use  $X$  to denote a generic random variable with the same distribution as all the  $X_t$ 's. Our focus is to assess the model misspecification effect when the identical probability distribution of the  $X_t$ 's deviates from  $P_0$ . The formulation in (1) is now written as

$$\begin{aligned} \max \quad & E_f[h(\mathbf{X}_T)] \\ \text{subject to} \quad & D(P_f||P_0) \leq \eta \\ & P_f \in \mathcal{P}_0. \end{aligned} \tag{7}$$

The function  $h(\cdot, \dots, \cdot)$  is now assumed to satisfy a majorization condition:

**Assumption 3.** The cost function  $h$  satisfies  $|h(\mathbf{X}_T)| \leq \sum_{t=1}^T \Lambda_t(X_t)$  a.s., for some deterministic functions  $\Lambda_t(\cdot)$ , where each of the  $\Lambda_t(X_t)$ 's possesses exponential moment, i.e.  $E_0[e^{\theta\Lambda_t(X)}] < \infty$  for  $\theta$  in a neighborhood of zero.

Assumption 3 is easy to check and holds in many contexts. In particular, it holds trivially when  $h$  is bounded. Some examples will be demonstrated in Section 7.

Next, we also assume a non-degeneracy condition that is analogous to Assumption 2; this requires introducing a function  $g(\cdot) := \mathcal{G}(h)(\cdot)$  where  $\mathcal{G}$  is a functional acted on  $h$  and  $g := \mathcal{G}(h)$  maps from  $\mathcal{X}$  to  $\mathbb{R}$ . This function  $g(x)$  is defined as the sum of individual conditional expectations of  $h(\mathbf{X}_T)$  over all time steps, i.e.

$$g(x) = \sum_{t=1}^T g_t(x) \quad (8)$$

where  $g_t(x)$  is the individual conditional expectation at time  $t$ , given by

$$g_t(x) = E_0[h(\mathbf{X}_T)|X_t = x]. \quad (9)$$

Our non-degeneracy condition is imposed on  $g$  acted on the individual random variable  $X$ :

**Assumption 4.** *The random variable  $g(X)$  is non-degenerate, i.e. non-constant, under the benchmark distribution  $P_0$ .*

Assumption 4 guarantees that  $P_0$  is not a “locally optimal” model, in the sense that there always exists a direction of model discrepancy that strictly dominates  $P_0$ . When  $T = 1$ , Assumption 4 is reduced to Assumption 2.

We have the following result:

**Theorem 2.** *With Assumptions 3 and 4 in hold, the optimal value of (7) satisfies*

$$E_{f^*}[h(\mathbf{X}_T)] = E_0[h(\mathbf{X}_T)] + \sqrt{2\text{Var}_0(g(X))}\eta^{1/2} + \frac{1}{\text{Var}_0(g(X))} \left( \frac{1}{3}\kappa_3(g(X)) + \nu \right) \eta + O(\eta^{3/2}) \quad (10)$$

where  $\text{Var}_0(g(X))$  and  $\kappa_3(g(X))$  are the variance and the third order cumulant of  $g(X)$  respectively, and

$$\nu = E_0[(G(X, Y) - E_0[G(X, Y)])(g(X) - E_0[g(X)])(g(Y) - E_0[g(Y)])]. \quad (11)$$

Here  $g(\cdot)$  is defined in (8), and  $G(\cdot, \cdot)$  is a function derived from  $h$  that is defined as

$$G(x, y) = \sum_{t=1}^T \sum_{s=1, \dots, T, s \neq t} G_{ts}(x, y) \quad (12)$$

where

$$G_{ts}(x, y) = E_0[h(\mathbf{X}_T)|X_t = x, X_s = y]. \quad (13)$$

Here  $X$  and  $Y$  are independent random variables each having distribution  $P_0$ .

The first and second order coefficients in (10) possess all the properties of nonparametric derivatives discussed in Section 3.1. Additionally, we note the following:

1. The function  $g(\cdot)$  can be interpreted as a “symmetrization” of the multivariate function  $h(\cdot, \dots, \cdot)$ , by collapsing the domain from  $\mathcal{X}^T$  to  $\mathcal{X}$  via sequential conditioning. This function  $g(\cdot)$  plays an important role in controlling the associated optimality conditions and comes up

naturally due to the i.i.d. assumptions on  $X_1, \dots, X_T$ , as will be seen in our development in Section 5. Besides, an alternate way to interpret  $g(\cdot)$  is to define a function  $S_h : \mathcal{X}^T \rightarrow \mathbb{R}$  as

$$\begin{aligned} S_h(x_1, \dots, x_T) \\ = h(x_1, x_2, \dots, x_T) + h(x_2, x_1, x_3, x_4, \dots, x_T) + h(x_3, x_2, x_1, x_4, \dots, x_T) + \dots \\ + h(x_T, x_2, x_3, \dots, x_1) \end{aligned} \quad (14)$$

i.e.  $S_h$  is the sum of all the swaps from the first coordinate with each of the other coordinates. Then  $g(x)$  is merely  $E_0[S_h(\mathbf{X}_T)|X_1 = x]$ . In the case that  $h$  is symmetric over all its arguments,  $S_h(\mathbf{X}_T)$  becomes  $Th(\mathbf{X}_T)$ , and  $g(x) = TE_0[h(\mathbf{X}_T)|X_1 = x]$ .

2. The function  $G(x, y)$  again has a “symmetrization” interpretation, by conditioning on two random components in each summand. Moreover, in the formula (11), one can replace  $G(x, y) = \sum_{t=1}^T \sum_{s=1, \dots, T, s \neq t} G_{ts}(x, y)$  by either

$$2 \sum_{t=1}^T \sum_{s < t} G_{ts}(x, y) \quad \text{or} \quad 2 \sum_{t=1}^T \sum_{s > t} G_{ts}(x, y)$$

by simple arithmetic and replacing the roles of  $X$  and  $Y$  for some summands. Furthermore, the formula (11) can be written as

$$\nu = E_0[G(X, Y)(g(X) - E_0[g(X)])(g(Y) - E_0[g(Y)])]$$

by using the independence of  $X$  and  $Y$ .

3. The first and second order nonparametric derivatives are translation invariant in both  $g(x)$  and  $G(x, y)$ , inherited from the translation invariance of cumulants and the centering for each factor in the expression of  $\nu$  in (11).

Theorem 2 is illustrated by the following example:

**Example 2** (Tail probability of i.i.d. sum). *Consider computing the probability  $P\left(\sum_{t=1}^T X_t > y\right)$  where  $X_t$  are i.i.d. Gaussian variables with mean 0 and variance  $\sigma^2$ . Suppose there is a change in the mean, within the Gaussian family, from 0 to  $\mu \neq 0$ . By the likelihood ratio method (see, for example, §7 in [3]), since  $(d/d\mu) \log \phi(x; \mu, \sigma^2)|_{\mu=0} = x/\sigma^2$ , where  $\phi(x; \mu, \sigma^2)$  denotes the density of  $N(\mu, \sigma^2)$ , the first order parametric derivative with respect to  $\mu$  is*

$$\begin{aligned} \left. \frac{dP\left(\sum_{t=1}^T X_t > y\right)}{d\mu} \right|_{\mu=0} &= E_0 \left[ I\left(\sum_{t=1}^T X_t > y\right) \sum_{t=1}^T \frac{X_t}{\sigma^2} \right] = \frac{E_0[\sum_{t=1}^T X_t; \sum_{t=1}^T X_t > y]}{\sigma^2} \\ &= \frac{E_0[N(0, T\sigma^2); N(0, T\sigma^2) > y]}{\sigma^2} \end{aligned}$$

To compare with our nonparametric framework, let us first convert the above calculation into a scale in terms of KL divergence. We know already from Example 1 that the KL divergence between  $N(0, \sigma^2)$  and  $N(\mu, \sigma^2)$  is  $\mu^2/(2\sigma^2)$ , which means a unit change in  $\mu$  is equivalent to  $\sqrt{2}\sigma$  change in the square root of KL divergence. So the first order derivative with respect to the square root of KL divergence, along this particular direction, is

$$\frac{\sqrt{2}}{\sigma} E_0[N(0, T\sigma^2); N(0, T\sigma^2) > y] = \sqrt{\frac{T}{\pi}} e^{-y^2/(2T\sigma^2)} \quad (15)$$

Now let us consider a change in the model beyond the Gaussian family, but still retaining the i.i.d. assumption. Note that Assumption 3 holds trivially since the cost function  $I(\sum_{t=1}^T X_t > y)$  is an indicator function and hence bounded. Next, we have  $g(x) = \sum_{t=1}^T P_0\left(\sum_{s=1}^T X_s > y | X_t = x\right) = TP_0\left(\sum_{s=1}^T X_s > y | X_1 = x\right)$  (alternatively, since  $I\left(\sum_{t=1}^T X_t > y\right)$  is symmetric, by discussion point 1 above,  $g(x) = TE_0[h(\mathbf{X}_T) | X_1 = x]$  giving the same conclusion). The random variable

$$g(X) = TP_0\left(X + \sum_{t=2}^T X_t > y \middle| X\right) = T\bar{\Phi}(y - X; 0, (T-1)\sigma^2)$$

where  $\bar{\Phi}(y; 0, \sigma^2)$  denotes the tail distribution of  $N(0, \sigma^2)$  and  $X \sim N(0, \sigma^2)$ , is clearly non-constant. Consequently, Assumption 4 is satisfied. Moreover, the first order nonparametric derivative for the tail probability is

$$\sqrt{2\text{Var}_0(g(X))} = T\sqrt{2\text{Var}_0(\bar{\Phi}(y - X; 0, (T-1)\sigma^2))}. \quad (16)$$

According to discussion point 6 in Section 3.1, the value of (16) must dominate (15). For example, when  $y = 10$ ,  $T = 5$ , and  $\sigma = 2$ , the parametric derivative with respect to the mean, given by (15), is 0.104. On the other hand, numerical estimation of the first order nonparametric derivative shown by (16) is 0.131 (within  $10^{-4}$  error with 95% confidence, using  $10^7$  samples). When  $y = 10$ ,  $T = 10$  and  $\sigma = 1$ , the parametric derivative (15) is 0.012 and the nonparametric derivative is 0.015 (within  $10^{-5}$  error with 95% confidence).

We caution that although a sum of Gaussian random variables  $X_1 + \dots + X_T$  is still Gaussian, which allows straightforward evaluation of the probability of interest as well as the parametric derivative along the movement in mean, a perturbation of model beyond the Gaussian family for the summand variables will destroy their infinite divisibility property [14]. In this sense there is no shortcut to obtain the formula (16).

Intuitively, in our nonparametric derivatives in Theorem 2, the moments of the sum of sequential conditional expectations, each given a particular time step or pairs of time steps, can be interpreted as a summary of the variation due to all the underlying individual components. The form of the first order nonparametric derivative stipulates that its estimation can suffer from a linear growth of mean square error with the time horizon. This issue resembles that of the likelihood ratio method in classical parametric derivative estimation. For certain performance measures, such as those that require tracking the system's large deviations properties, it is plausible that variance reduction techniques can be employed, as in the parametric case [41]. Moreover, stationarity can potentially play a role in achieving an asymptotic mean square error of the estimators that is sublinear in  $T$ .

### 3.3 Random Time Horizon Problems

We now extend our result to problems involving a random time  $\tau$ . Consider the cost function  $h(\mathbf{X}_\tau)$  that depends on the sequence  $\mathbf{X}_\tau = (X_1, X_2, \dots, X_\tau)$ . Our formulation is then

$$\begin{aligned} & \max && E_f[h(\mathbf{X}_\tau)] \\ & \text{subject to} && D(P_f \| P_0) \leq \eta \\ & && P_f \in \mathcal{P}_0 \end{aligned} \quad (17)$$

Let us lay out our assumptions on  $h$  and the random time  $\tau$ . We impose either a boundedness or an independence condition on  $\tau$ :

**Assumption 5.** The random time  $\tau$  is a stopping time with respect to  $\{\mathcal{F}_t\}_{t \geq 1}$ , a filtration that supersedes the filtration that is generated by the sequence  $\{X_t\}_{t \geq 1}$ , namely  $\{\mathcal{F}(X_1, \dots, X_t)\}_{t \geq 1}$ . Moreover,  $\tau$  is bounded a.s. by a deterministic time  $T$ . The cost function  $h$  satisfies  $|h(\mathbf{X}_\tau)| \leq \sum_{t=1}^T \Lambda_t(X_t)$  a.s. for some deterministic functions  $\Lambda_t(\cdot)$ , where  $\Lambda_t(X_t)$  each possesses finite exponential moment, i.e.  $E_0[e^{\theta \Lambda_t(X)}] < \infty$ , for  $\theta$  in a neighborhood of zero.

**Assumption 6.** The random time  $\tau$  is independent of the sequence  $\{X_t\}_{t \geq 1}$ , and has finite second moment under  $P_0$ , i.e.  $E_0\tau^2 < \infty$ . Moreover, the cost function  $h(\mathbf{X}_\tau)$  is bounded a.s..

Next, we also place a non-degeneracy condition analogous to Assumption 4. With  $\tau$  satisfying either of the assumptions above, we define  $\tilde{g} : \mathcal{X} \rightarrow \mathbb{R}$  as

$$\tilde{g}(x) = \sum_{t=1}^{\infty} \tilde{g}_t(x) \quad (18)$$

where  $\tilde{g}_t(x)$  is given by

$$\tilde{g}_t(x) = E_0[h(\mathbf{X}_\tau); \tau \geq t | X_t = x]. \quad (19)$$

Our non-degeneracy condition is now imposed on the function  $\tilde{g}$  acted on  $X$ :

**Assumption 7.** The random variable  $\tilde{g}(X)$  is non-degenerate, i.e. non-constant, under the benchmark distribution  $P_0$ .

Either one of Assumption 5 or 6 suffices to extend our scheme in Section 3.2 to random time horizon. We have the following theorem:

**Theorem 3.** With either Assumption 5 or 6 in hold, together with Assumption 7, the optimal value of (17) satisfies

$$E_{f^*}[h(\mathbf{X}_\tau)] = E_0[h(\mathbf{X}_\tau)] + \sqrt{2\text{Var}_0(\tilde{g}(X))}\eta^{1/2} + \frac{1}{\text{Var}_0(\tilde{g}(X))} \left( \frac{1}{3}\kappa_3(\tilde{g}(X)) + \tilde{\nu} \right) \eta + O(\eta^{3/2}) \quad (20)$$

where

$$\tilde{\nu} = E_0[(\tilde{G}(X, Y) - E_0[\tilde{G}(X, Y)])(\tilde{g}(X) - E_0[\tilde{g}(X)])(\tilde{g}(Y) - E_0[\tilde{g}(Y)])]. \quad (21)$$

Here  $\tilde{g}(x)$  is defined in (18), and  $\tilde{G}(x, y)$  is defined as  $\tilde{G}(x, y) = \sum_{t=1}^{\infty} \sum_{\substack{s \geq 1 \\ s \neq t}} \tilde{G}_{ts}(x, y)$ , where

$\tilde{G}_{ts}(x, y)$  is given by

$$\tilde{G}_{ts}(x) = E_0[h(\mathbf{X}_\tau); \tau \geq t \wedge r | X_t = x, X_s = y].$$

To get some intuition, note that the function  $g(x)$  introduced in Theorem 2 in the last subsection can be expressed as

$$g(x) = \sum_{t=1}^T E_0[h(\mathbf{X}_T) | X_t = x] = \sum_{t=1}^{\infty} E_0[h(\mathbf{X}_T) | X_t = x] I(T \geq t) = \sum_{t=1}^{\infty} E_0[h(\mathbf{X}_T); T \geq t | X_t = x]. \quad (22)$$

Of course, this calculation is only valid when  $T$  is deterministic. Theorem 3, however, states that we can replace  $T$  in (22) by a random time  $\tau$  under the proper assumptions.

Similar to Theorem 2, the function  $\tilde{G}(x, y)$  in (21) can be replaced by

$$2 \sum_{t=1}^{\infty} \sum_{s < t} E_0[h(\mathbf{X}_\tau); \tau \geq s | X_t = x, X_s = y] \text{ or } 2 \sum_{t=1}^{\infty} \sum_{s > t} E_0[h(\mathbf{X}_\tau); \tau \geq t | X_t = x, X_s = y].$$

Regarding computation, especially using Monte Carlo method, the fact that  $\tilde{g}(x)$  and  $\tilde{G}(x, y)$  involve infinite sums may call for some anxiety. One way to get around is to express the sums as expectations over infinitely supported random variables along with appropriate weighting. For example, in the case of  $\tilde{g}(x)$ , we can introduce a random time  $R$ , independent of the system, with probability mass function  $p_\omega, \omega = 1, 2, \dots$ . Then

$$\tilde{g}(x) = E_0[p_R^{-1} h(\mathbf{X}_\tau^R); \tau \geq R | X = x]$$

where  $\mathbf{X}_\tau^R$  is  $\mathbf{X}_\tau$  with the  $R$ -th component replaced by  $X = x$ . In other words, one can simulate  $\tilde{g}(x)$  by the following procedure (with  $x$  fixed):

1. Sample the auxiliary random time  $R$ ; say  $R = \omega$  is the realization.
2. Sample  $X_1, \dots, X_\tau$ , with  $X_\omega$  fixed as  $x$ .
3. If  $\tau < \omega$ , output 0; else output  $p_\omega^{-1} h(\mathbf{X}_\tau)$ .

This yields a sample of  $p_R^{-1} h(\mathbf{X}_\tau^R) I(\tau \geq R)$  given  $X = x$ . The choice of the random time  $R$  can be quite flexible, as long as the involved expectation (and also second moment for the sake of consistency) is finite. This time horizon randomization technique has been used in devising importance samplers for first passage and steady-state problems in queueing networks and Gaussian processes (see, for example, [9, 8]).

### 3.4 Auxiliary Variables

In some applications, the cost function can involve many different random sources, and one is interested in evaluating the model misspecification effect from one particular source. Our framework discussed in the last three subsections can be easily adopted to such scenarios. Suppose there is a random object  $\mathbf{Y}$  in the system, potentially dependent of  $\{X_t\}_{t \geq 1}$ , and we are interested in the model misspecification effect of the distribution of  $X$  for the performance measure  $E_0[h(\mathbf{X}, \mathbf{Y})]$ . This random object  $\mathbf{Y}$  can be a sequence of random variables, say  $Y_1, Y_2, \dots$ , or can be in other plausible forms. All the results discussed in the previous subsections, namely Theorems 1, 2 and 3, still hold, but replacing  $h(\mathbf{X})$  by  $E_0[h(\mathbf{X}, \mathbf{Y}) | \mathbf{X}]$  (Part of the statements in the theorems can then be further simplified). Such a direct replacement is enough because one can consider  $E_f[E_0[h(\mathbf{X}, \mathbf{Y}) | \mathbf{X}]]$  as the performance measure and  $E_0[h(\mathbf{X}, \mathbf{Y}) | \mathbf{X}]$  as the cost function, noting that  $E_0$  here involves only the random object  $\mathbf{Y}$  which is not subject to perturbation.

## 4 Explanation of the Preliminary Results

In this and the following section, we will explain the main methodology in arriving at the theorems in Section 3. This section will be devoted to the basic case, namely Theorem 1, which will also serve as an important building block for the other results.

To solve the maximization problem (4), we first transform the decision variables from the space of measures to the space of functions, which is easier to handle. Recall that  $P_f$  is assumed to be

absolutely continuous with respect to  $P_0$ , and hence the likelihood ratio  $L := dP_f/dP_0$  exists. Via a change of measure, the optimization problem (4) can be rewritten as a maximization over the likelihood ratios, i.e. (4) is equivalent to

$$\begin{aligned} & \max && E_0[h(X)L(X)] \\ & \text{subject to} && E_0[L(X) \log L(X)] \leq \eta \\ & && L \in \mathcal{L} \end{aligned} \quad (23)$$

where  $\mathcal{L} := \{L \in \mathcal{L}_1(P_0) : E_0[L] = 1, L \geq 0 \text{ a.s.}\}$ , and we denote  $\mathcal{L}_1(P_0)$  as the  $\mathcal{L}_1$ -space with respect to the measure  $P_0$  (we sometimes suppress the dependence of  $X$  in  $L = L(X)$  for convenience when no confusion arises). The key then is to find an optimal solution  $L^*$ , and investigate its asymptotic relation with  $\eta$ .

To solve (23), we consider the Lagrangian relaxation

$$\min_{\alpha \geq 0} \max_{L \in \mathcal{L}} E_0[h(X)L] - \alpha(E_0[L \log L] - \eta). \quad (24)$$

Since the constraint is convex and the objective is linear, strong duality holds [39], and the optimal values of (23) and (24) coincide. Consider the inner maximization of (24) given by

$$\max_{L \in \mathcal{L}} E_0[h(X)L - \alpha L \log L]. \quad (25)$$

The solution for this inner maximization is characterized by the following proposition:

**Proposition 1.** *Suppose Assumption 1 holds, and consider  $\alpha > 0$ . When  $\alpha$  is sufficiently large, there exists a unique optimizer of (25) given by*

$$L^*(x) = \frac{e^{h(x)/\alpha}}{E_0[e^{h(X)/\alpha}]} \quad (26)$$

This result is known in robust single-stage control [27]. Let us review briefly here. A heuristic for getting (26) is to apply Euler-Lagrange equation and informally differentiate the integrand with respect to  $L$ . First, the Lagrangian relaxation of the constraint  $E_0[L] = 1$  is

$$E_0[h(X)L - \alpha L \log L + \lambda L - \lambda]$$

where  $\lambda \in \mathbb{R}$  is the Lagrange multiplier. Treating  $E_0[\cdot]$  as an integral, Euler-Lagrange equation implies that the derivative with respect to  $L$  is

$$h(X) - \alpha \log L - \alpha + \lambda = 0$$

which gives

$$\log L = \frac{h(X)}{\alpha} + \frac{\lambda - \alpha}{\alpha}$$

or that  $L = \lambda e^{h(X)/\alpha}$  for some  $\lambda > 0$ . With the constraint that  $E_0[L] = 1$ , a candidate solution is

$$L^* = \frac{e^{h(X)/\alpha}}{E_0[e^{h(X)/\alpha}]} \quad (27)$$

The above argument is heuristic. To verify (27) formally, the following convexity argument will suffice. First, note that the objective value of (25) evaluated at  $L^*$ , given by (27), is

$$\begin{aligned} E_0[h(X)L^* - \alpha L^* \log L^*] &= E_0 \left[ h(X)L^* - \alpha L^* \left( \frac{h(X)}{\alpha} - \log E_0[e^{h(X)/\alpha}] \right) \right] \\ &= \alpha \log E_0[e^{h(X)/\alpha}] \end{aligned} \quad (28)$$

Our goal is to show that

$$\alpha \log E_0[e^{h(X)/\alpha}] \geq E_0[h(X)L - \alpha L \log L]$$

for all  $L \in \mathcal{L}$ . Rearranging terms, this means we need

$$E_0[e^{h(X)/\alpha}] \geq e^{E_0[h(X)L - \alpha L \log L]/\alpha} \quad (29)$$

To prove (29), observe that, for any likelihood ratio  $L$ ,

$$E_0[e^{h(X)/\alpha}] = E_0[LL^{-1}e^{h(X)/\alpha}] = E_0[Le^{h(X)/\alpha - \log L}] \geq e^{E_0[h(X)L/\alpha - L \log L]}$$

by using the convexity of the function  $e^{\cdot}$  and Jensen's inequality over the expectation  $E_0[L \cdot]$  in the last inequality. Note that equality holds if and only if  $h(X)/\alpha - \log L$  is degenerate, i.e.  $h(X)/\alpha - \log L = \text{constant}$ , which reduces to  $L^*$ . Hence uniqueness also holds.

In conclusion, when  $1/\alpha \in \mathcal{D}^+ := \{\theta \in \mathbb{R}^+ \setminus \{0\} : \psi(\theta) < \infty\}$ , the optimal solution of (25) is given by (26), with the optimal value  $\alpha \log E_0[e^{h(X)/\alpha}]$ . This fact will be used in the proof of Theorem 1:

*Proof of Theorem 1.* We want to express the optimal value of (24) in terms of  $\eta$ . Let  $\alpha^*$  be the optimal dual solution. Our proof scheme is divided into the following two steps:

**Relation between  $\eta$  and  $\alpha^*$ .** By complementary slackness, either  $\alpha^* = 0$  or, in the case  $\alpha^* > 0$ ,  $E_0[L^* \log L^*] = \eta$ . Let us assume, for the moment, that the latter case holds, i.e.  $\alpha^* > 0$ . Moreover, we assume that  $\alpha^*$  is sufficiently large, namely that  $1/\alpha^* \in \mathcal{D}^+$ . We will exclude the other possibilities on  $\alpha^*$  at the end of this proof. Now we have

$$\begin{aligned} \eta = E_0[L^* \log L^*] &= \frac{E[h(X)L^*]}{\alpha^*} - \log E_0[e^{h(X)/\alpha^*}] \\ &= \frac{\beta E_0[h(X)e^{\beta h(X)}]}{E_0[e^{\beta h(X)}]} - \log E_0[e^{\beta h(X)}] = \beta \psi'(\beta) - \psi(\beta) \end{aligned} \quad (30)$$

where  $\beta = 1/\alpha^*$ , and  $\psi(\beta) = \log E_0[e^{\beta h(X)}]$  is the logarithmic moment generating function of  $h(X)$ .

**Relation between the optimal primal objective value and  $\alpha^*$ .** The primal objective is

$$E_0[h(X)L^*] = \frac{E_0[h(X)e^{h(X)/\alpha^*}]}{E_0[e^{h(X)/\alpha^*}]} = \frac{E_0[h(X)e^{\beta h(X)}]}{E_0[e^{\beta h(X)}]} = \psi'(\beta) \quad (31)$$

This gives the first part of the theorem. It is then a simple application of Taylor's expansion to arrive at the asymptotic relation in (6). First, note that the relation (30) can be written as

$$\begin{aligned}
\eta &= \beta\psi'(\beta) - \psi(\beta) = \sum_{n=0}^{\infty} \frac{1}{n!} \kappa_{n+1} \beta^{n+1} - \sum_{n=0}^{\infty} \frac{1}{n!} \kappa_n \beta^n \\
&= \sum_{n=1}^{\infty} \left[ \frac{1}{(n-1)!} - \frac{1}{n!} \right] \kappa_n \beta^n = \sum_{n=2}^{\infty} \frac{1}{n(n-2)!} \kappa_n \beta^n \\
&= \frac{1}{2} \kappa_2 \beta^2 + \frac{1}{3} \kappa_3 \beta^3 + \frac{1}{8} \kappa_4 \beta^4 + O(\beta^5)
\end{aligned} \tag{32}$$

where  $\kappa_n = \psi^{(n)}(0)$  is the  $n$ -th cumulant of  $h(X)$ . By Assumptions 1 and 2,  $\psi(\cdot)$  is strictly convex, so  $\beta\psi'(\beta) - \psi(\beta)$  has a unique zero at  $\beta = 0$ . Consequently, we can invert the relation (32) to get

$$\begin{aligned}
\beta &= \sqrt{\frac{2\eta}{\kappa_2}} \left( 1 + \frac{2}{3} \frac{\kappa_3}{\kappa_2} \beta + \frac{1}{4} \frac{\kappa_4}{\kappa_2} \beta^2 + O(\beta^3) \right)^{-1/2} \\
&= \sqrt{\frac{2\eta}{\kappa_2}} \left( 1 - \frac{1}{3} \frac{\kappa_3}{\kappa_2} \beta + O(\beta^2) \right) \\
&= \sqrt{\frac{2}{\kappa_2}} \eta^{1/2} - \frac{2}{3} \frac{\kappa_3}{\kappa_2^2} \eta + O(\eta^{3/2})
\end{aligned}$$

As a result, (31) can be expanded as

$$\begin{aligned}
E_0[h(X)L^*] &= \psi'(\beta) = \kappa_1 + \kappa_2 \beta + \kappa_3 \frac{\beta^2}{2} + O(\beta^3) \\
&= \kappa_1 + \kappa_2 \left( \sqrt{\frac{2}{\kappa_2}} \eta^{1/2} - \frac{2}{3} \frac{\kappa_3}{\kappa_2^2} \eta + O(\eta^{3/2}) \right) + \frac{\kappa_3}{2} \left( \frac{2}{\kappa_2} \eta + O(\eta^{3/2}) \right) + O(\eta^{3/2}) \\
&= \kappa_1 + \sqrt{2\kappa_2} \eta^{1/2} + \frac{1}{3} \frac{\kappa_3}{\kappa_2} \eta + O(\eta^{3/2})
\end{aligned}$$

which gives (6).

To conclude the theorem, we need to argue that  $\alpha^*$  is indeed positive and is sufficiently large, namely that  $1/\alpha^* \in \mathcal{D}^+$ . Since the objective of the dual problem, i.e. the objective in the outer minimization in (24), is convex, we need only show that our choice of  $\alpha^*$  that satisfies (30) is a local minimum. To this end, note that any  $\alpha$  in a neighborhood of  $\alpha^*$  has dual objective function  $\alpha\psi(1/\alpha) + \alpha\eta$  (which can be seen by, for example, the calculation in (28)). The derivative of this objective function  $(d/d\alpha)(\alpha\psi(1/\alpha) + \alpha\eta) = \psi(1/\alpha) - (1/\alpha)\psi'(1/\alpha) + \eta$  is equal to 0 at  $\alpha^*$ . Moreover, it is negative before and positive after  $\alpha^*$  in its immediate neighborhood, by a simple analysis using the strict convexity of  $\psi(\cdot)$ . This concludes that  $\alpha^*$  chosen in (30) must be the unique optimal dual solution.  $\square$

The simple convexity argument to obtain  $L^*$  in Proposition 1 no longer suffices for multiple time horizon. In that situation, the structure of the optimal change of measure will not be a simple exponential tilting, but instead will be characterized as the fixed point of a contraction map on the  $\mathcal{L}_1$ -space. The next section will focus on this development.

## 5 Main Methodology

Our goal in this section is to lay out the methodology in proving the result for general time horizon  $T > 1$ , namely Theorem 2. The cost function  $h(\cdot, \dots, \cdot)$  now depends on an i.i.d. sequence  $\{X_t\}_{t=1, \dots, T}$ , each with benchmark distribution  $P_0$ . Leveraging the idea in the previous section, we first write the maximization problem (7) in terms of likelihood ratio  $L$ :

$$\begin{aligned} & \max && E_0[h(\mathbf{X}_T)\underline{L}_T] \\ & \text{subject to} && E_0[L(X) \log L(X)] \leq \eta \\ & && L \in \mathcal{L} \end{aligned} \tag{33}$$

where for convenience we denote  $\underline{L}_T = \prod_{t=1}^T L(X_t)$ , and  $X$  as a generic variable that is independent of  $\{X_t\}_{t=1, \dots, T}$  and having identical distribution as each  $X_t$ . The following is the recipe from the last section that we will follow:

1. Consider the Lagrangian relaxation of (33), and characterize the solution for the inner maximization.
2. Argue that the Lagrangian relaxation gives the optimal value for the primal objective in (33).
3. Expand the solution in the dual variable  $\alpha^*$  and hence  $\eta$ .

For  $T > 1$ , the main technical challenge is the product form  $\underline{L}_T$  that appears in the objective function in (33). This hinders a direct convexity argument in finding an optimal change of measure  $L^*$ . To get around, we shall introduce a fixed point equation to find a candidate solution, and use an induction argument to establish optimality. The following subsections will implement these step-by-step.

### 5.1 Optimal Change of Measure via Fixed Point Equation

Before we start our analysis, let us first discuss some features of the optimization problem (33). As a comparison, suppose that we treat  $\mathbf{X}_T$  as a single variable, and consider a deviation of KL divergence within  $T\eta$  units. Then the optimization formulation following (23) in Section 4 will give

$$\begin{aligned} & \max && E_0[h(\mathbf{X}_T)L(\mathbf{X}_T)] \\ & \text{subject to} && E_0[L(\mathbf{X}_T) \log L(\mathbf{X}_T)] \leq T\eta \\ & && L \in \mathcal{L}^T \end{aligned} \tag{34}$$

where  $\mathcal{L}^T$  is now defined on the  $T$ -product measure of  $P_0$ . This primitive formulation (34) provides an upper bound on the optimal value of (33). It can be seen by writing the constraint in (33) as  $E[\underline{L}_T \log \underline{L}_T] \leq T\eta$ , since  $X_t$ 's are i.i.d., and then relaxing the implicit constraint that  $\underline{L}_T = \prod_{t=1}^T L(X_t)$ .

This observation also implies that (33) has a bounded optimal value, i.e. it is bounded by the optimal value of (34). The latter is finite because Assumption 3 implies that

$$E_0[e^{\theta h(\mathbf{X}_T)}] \leq E_0[e^{\theta \sum_{t=1}^T \Lambda_t(X_t)}] \leq \prod_{t=1}^T E_0[e^{\theta \Lambda_t(X_t)}] < \infty \tag{35}$$

when  $\theta$  is small enough. Hence  $h(\mathbf{X}_T)$  has exponential moment and Assumption 1 is satisfied. It is also trivial to check that Assumption 4 implies Assumption 2. Therefore, the result in Theorem 1 applies to (34).

Now, focusing back on the formulation (33), we shall proceed by first reducing the size of the space that  $L$  subsists in the formulation. We will soon see that this is useful for constructing an associated contraction map. Moreover, we shall define a metric for the contraction to facilitate our later development of monotonicity properties in reaching the involved objective values. Let  $\Lambda(x) = \sum_{t=1}^T \Lambda_t(x)$  where  $\Lambda_t(x)$  is defined in Assumption 3. Define

$$\mathcal{L}(M) = \{L \in \mathcal{L} : E_0[\Lambda(X)L(X)] \leq M\}$$

for  $M > 0$ , and the associated norm  $\|L\|_\Lambda := E_0[(1 + \Lambda(X))L(X)]$  and hence metric

$$\|L - L'\|_\Lambda = E_0[(1 + \Lambda(X))|L(X) - L'(X)|]. \quad (36)$$

It is routine to check that  $\mathcal{L}(M)$  is complete. The following observation is useful for our development:

**Lemma 1.** *For any  $\eta \leq N$  for some small  $N$ , the formulation (33) is equivalent to*

$$\begin{aligned} & \max && E_0[h(\mathbf{X}_T)\underline{L}_T] \\ & \text{subject to} && E_0[L(X) \log L(X)] \leq \eta \\ & && L \in \mathcal{L}(M) \end{aligned} \quad (37)$$

for some large enough  $M > 0$ , independent of  $\eta$ .

*Proof.* As in the rest of this paper, we shall suppress the dependence of  $X$  in  $L = L(X)$  whenever no confusion arises. We want to show that  $E_0[L \log L] \leq \eta$  and  $L \in \mathcal{L}$  together implies  $L \in \mathcal{L}(M)$ . Note that  $\Lambda(X)$  has exponential moment, since Holder's inequality implies

$$E_0[e^{\theta \Lambda(X)}] = E_0[e^{\theta \sum_{t=1}^T \Lambda_t(X)}] \leq \prod_{t=1}^T (E_0[e^{T\theta \Lambda_t(X)}])^{1/T} < \infty \quad (38)$$

when  $\theta$  is small enough. Hence, for any  $L \in \mathcal{L}$  that satisfies  $E_0[L \log L] \leq \eta$ , we have

$$E_0[e^{\theta \Lambda(X)}] = E_0[LL^{-1}e^{\theta \Lambda(X)}] = E_0[Le^{\theta \Lambda(X) - \log L}] < \infty$$

for small enough  $\theta$ , by (38). Jensen's inequality implies that

$$e^{\theta E_0[\Lambda(X)L] - E_0[L \log L]} \leq E_0[Le^{\theta \Lambda(X) - \log L}] = E_0[e^{\theta \Lambda(X)}] < \infty.$$

Since  $E_0[L \log L] \leq \eta \leq N$ , we have  $E_0[\Lambda(X)L] \leq M$  for some constant  $M$ . So  $L \in \mathcal{L}(M)$ . This concludes the lemma.  $\square$

The Lagrangian relaxation of (37) is given by

$$\min_{\alpha \geq 0} \max_{L \in \mathcal{L}(M)} E_0[h(\mathbf{X}_T)\underline{L}_T] - \alpha(E_0[L \log L] - \eta) \quad (39)$$

with inner maximization

$$\max_{L \in \mathcal{L}(M)} E_0[h(\mathbf{X}_T)\underline{L}_T] - \alpha E_0[L \log L]. \quad (40)$$

Note that the first expectation  $E_0[\cdot]$  in (40) is acted on the product space of  $\{X_t\}_{t=1, \dots, T}$ . To get some intuition on how to solve (40), note that when  $h$  is separable and symmetric, i.e.  $h(\mathbf{X}_T) = \sum_{t=1}^T w(X_t)$  for some  $w(\cdot)$ , the quantity  $E_0[h(\mathbf{X}_T)\underline{L}_T]$  is also separable as  $E_0[h(\mathbf{X}_T)\underline{L}_T] = \sum_{t=1}^T E_0[w(X_t)L(X_t)] = TE_0[w(X)L(X)]$ . So the solution for (40) reduces to Proposition 1 but

with  $\alpha$  replaced by  $\alpha/T$ . The challenge is the general case when such separation for  $h$  does not hold. In this case, one useful way to view  $E_0[h(\mathbf{X}_T)\underline{L}_T]$ , as a mimic of the aforementioned separability, is through writing

$$TE_0[h(\mathbf{X}_T)\underline{L}_T] = \sum_{t=1}^T E_0[g_t^L(X_t)L(X_t)] = E_0[g^L(X)L(X)]$$

where  $g_t^L(x) = E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L(X_r) \middle| X_t = x \right]$  and  $g^L(x) = \sum_{t=1}^T g_t^L(x)$ . The functions  $g_t^L$  and  $g^L$  will appear frequently in the sequel. Note that  $g^L$  now depends on  $L$ , which differs from the symmetric separable case.

The analog of Proposition 1 for time horizon  $T > 1$  is:

**Proposition 2.** *Suppose Assumption 3 holds. When  $\alpha > 0$  is large enough, the unique optimal solution of (40) satisfies*

$$L(x) = \frac{e^{g^L(x)/\alpha}}{E_0[e^{g^L(X)/\alpha}]} \quad (41)$$

where  $g^L(x) = \sum_{t=1}^T g_t^L(x)$  and  $g_t^L(x) = E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L(X_r) \middle| X_t = x \right]$ .

Note that the function  $g^L(x)$  can be viewed as the expectation of the function  $S_h$  defined in (14), under the shifted measure  $P_f$ . In other words,

$$g^L(x) = \sum_{t=1}^T E_f[h(\mathbf{X}_T)|X_t = x] = E_f[S_h(\mathbf{X}_T)|X_1 = x].$$

As in Section 4, the form in (41) can be guessed from a heuristic differentiation with respect to  $L$ . Consider the Lagrangian relaxation of the constraint  $E_0[L] = 1$  in (40):

$$E_0[h(\mathbf{X}_T)L(X_1)L(X_2)\cdots L(X_T)] - \alpha E_0[L \log L] + \lambda E_0[L] - \lambda \quad (42)$$

There are  $T$  factors of  $L$  in the first term. A heuristic “product rule” is to differentiate each  $L$  factor, keeping all other  $L$ ’s unchanged. To do so, we condition on  $X_t$  to write

$$E_0[h(\mathbf{X}_T)L(X_1)L(X_2)\cdots L(X_T)] = E_0 \left[ E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L(X_r) \middle| X_t \right] L(X_t) \right]$$

and it turns out that the “right” differentiation of  $L$  on this quantity is represented as a sum of conditional expectations

$$\frac{d}{dL(x)} E_0 \left[ h(\mathbf{X}_T) \prod_{t=1}^T L(X_t) \right] \text{ “=” } \sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L(X_r) \middle| X_t = x \right]. \quad (43)$$

So the “product rule” gives an Euler-Lagrange equation as

$$\sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L(X_r) \middle| X_t = x \right] - \alpha \log L(x) - \alpha + \lambda = 0 \quad (44)$$

which gives

$$L(x) \propto \exp \left\{ \sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L(X_r) \middle| X_t = x \right] \right\} / \alpha$$

The constraint  $E_0[L] = 1$  then enforces the expression (41). To gain more understanding of the “product rule” (43), let us illustrate with a finitely supported  $X$ :

**Example 3.** Consider two i.i.d. random variables  $X_1$  and  $X_2$ , and a cost function  $h(X_1, X_2)$ . The variables  $X_1$  and  $X_2$  have finite support on  $1, 2, \dots, n$  under  $P_0$ . Denote  $p(x) = P_0(X_1 = x)$  for  $x = 1, 2, \dots, n$ . The objective value in (33) in this case is

$$E_0[h(X_1, X_2)L(X_1)L(X_2)] = \sum_{x_1=1}^n \sum_{x_2=1}^n h(x_1, x_2)p(x_1)p(x_2)L(x_1)L(x_2)$$

Now differentiate with respect to each  $L(1), L(2), \dots, L(n)$  respectively. For  $i = 1, \dots, n$ , we have

$$\begin{aligned} \frac{d}{dL(i)} E_0[h(X_1, X_2)] &= \sum_{x_1 \neq i} h(x_1, i)p(x_1)p(i)L(x_1) + \sum_{x_2 \neq i} h(i, x_2)p(i)p(x_2)L(x_2) + 2h(i, i)p(i)^2L(i) \\ &= E_0[h(X_1, i)L(X_1)] + E_0[h(i, X_2)L(X_2)]. \end{aligned}$$

This coincides with the product rule (43) discussed above.

We shall justify the above heuristic rigorously. The case for  $T = 2$  will be presented in the next subsection. We will then outline the general case, whose idea is similar but involves a few extra technical constructions. The full proof of the general case will be deferred to the appendix.

## 5.2 Proof of Proposition 2 for $T = 2$

In the case  $T = 2$ ,  $g^L(x) = E_0[h(X_1, X_2)L(X_2)|X_1 = x] + E_0[h(X_1, X_2)L(X_1)|X_2 = x] = E_0[S_h(X_1, X_2)L(X_2)|X_1 = x]$  where  $S_h$  is defined in (14). The proof of Proposition 2 centers around the operator  $\mathcal{K} := \mathcal{K}(h, \alpha)(\cdot) : \mathcal{L}(M) \rightarrow \mathcal{L}(M)$ , defined as

$$\mathcal{K}(L)(x) = \frac{e^{g^L(x)/\alpha}}{E_0[e^{g^L(X)/\alpha}]} = \frac{e^{E_0[S_h(X_1, X_2)L(X_2)|X_1=x]/\alpha}}{E_0[e^{E_0[S_h(X_1, X_2)L(X_2)|X_1]/\alpha}]} \quad (45)$$

Our argument is mainly divided into three steps. First, we prove that  $\mathcal{K}$  is a well-defined contraction mapping on  $\mathcal{L}(M)$ . This implies the existence and uniqueness of a fixed point in  $\mathcal{L}(M)$ . In the second step, we show that the objective value of the inner maximization (40) is, in certain sense, monotone in the iterations of  $\mathcal{K}$ . The third step comprises proving that this objective value converges to the value evaluated at the fixed point, which will then conclude Proposition 2.

For convenience, in the subsequent calculation we let  $C$  be a positive constant, not necessarily the same every time it appears.

**Step 1: Contraction Mapping.** We prove that  $\mathcal{K}$  is a well-defined contraction map on  $\mathcal{L}(M)$ , equipped with the norm defined in (36). The existence and uniqueness of a fixed point for  $\mathcal{K}$  is established by Banach fixed point theorem (§1 in [1]).

**Lemma 2.** *With Assumption 3, for sufficiently large  $\alpha$ , the operator  $\mathcal{K} : \mathcal{L}(M) \rightarrow \mathcal{L}(M)$  is well-defined, closed, and a strict contraction in  $\mathcal{L}(M)$  equipped with the norm  $\|\cdot\|_\Lambda$  introduced in (36). Hence there exists a unique fixed point  $L^* \in \mathcal{L}(M)$  that satisfies  $\mathcal{K}(L^*) = L^*$ .*

*Proof.* We show each element in the lemma one by one:

Well-definedness: Let  $\beta = 1/\alpha$  for convenience. To prove that  $\mathcal{K}$  is well-defined on  $\mathcal{L}(M)$ , we shall show that, for any  $L \in \mathcal{L}(M)$ ,  $0 < E_0[e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}] < \infty$  for small enough  $\beta$ . Recall Assumption 3 and the notation  $\Lambda(x) = \Lambda_1(x) + \Lambda_2(x)$  in (38). We have

$$\begin{aligned} E_0[e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}] &\leq E_0[e^{\beta E_0[(\Lambda(X_1) + \Lambda(X_2))L(X_2)|X_1]}] \\ &= E_0[e^{\beta(\Lambda(X_1) + E_0[\Lambda(X_2)L(X_2)])}] \\ &= E_0[e^{\beta\Lambda(X)}]e^{\beta E_0[\Lambda(X)L(X)]} \\ &\leq E_0[e^{\beta\Lambda(X)}]e^{\beta M} \\ &< \infty \end{aligned}$$

for  $L \in \mathcal{L}(M)$  and small enough  $\beta$ , where we use the independence of  $X_1$  and  $X_2$  in the first equality, the definition of  $\mathcal{L}(M)$  in the second inequality and (38) in the last inequality. Similarly, we also have

$$E_0[e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}] \geq E_0[e^{-\beta E_0[(\Lambda(X_1) + \Lambda(X_2))L(X_2)|X_1]}] \geq E_0[e^{-\beta\Lambda(X)}]e^{-\beta M} > 0 \quad (46)$$

for small enough  $\beta$ . Hence the denominator in the definition of  $\mathcal{K}$  is finite and non-zero. As a consequence, we have also shown that the numerator of  $\mathcal{K}$  given by  $e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1=x]}$  is finite a.s.. Therefore  $\mathcal{K}$  is well-defined.

Closedness: Next we show that  $\mathcal{K}$  is closed in  $\mathcal{L}(M)$  for large enough  $M$ , or more precisely, when  $M > E_0[\Lambda(X)]$ . For any  $L \in \mathcal{L}(M)$ , we have, by similar argument as above, that

$$\begin{aligned} E_0[\Lambda(X)\mathcal{K}(L)(X)] &= \frac{E_0[\Lambda(X_1)e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}]}{E_0[e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}]} \leq \frac{E_0[\Lambda(X)e^{\beta\Lambda(X)}]e^{2\beta E_0[\Lambda(X)L(X)]}}{E_0[e^{-\beta\Lambda(X)}]} \\ &\leq \frac{E_0[\Lambda(X)e^{\beta\Lambda(X)}]e^{2\beta M}}{E_0[e^{-\beta\Lambda(X)}]} \end{aligned} \quad (47)$$

As  $\beta \rightarrow 0$ , we have  $E_0[\Lambda(X)e^{\beta\Lambda(X)}] \rightarrow E_0[\Lambda(X)]$  and  $E_0[e^{-\beta\Lambda(X)}] \rightarrow 1$ . Hence, by choosing  $M > E_0[\Lambda(X)]$ , (47) is bounded by  $M$  when  $\beta$  is small enough. The result then follows.

Contraction: To check that  $\mathcal{K}$  is a contraction, consider, for any  $L, L' \in \mathcal{L}(M)$ ,

$$\begin{aligned}
\|\mathcal{K}(L) - \mathcal{K}(L')\|_\Lambda &= E_0[(1 + \Lambda(X))|\mathcal{K}(L)(X) - \mathcal{K}(L')(X)|] \\
&= E_0 \left[ (1 + \Lambda(X_1)) \left| \frac{e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}}{E_0[e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}]} - \frac{e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}}{E_0[e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}]} \right| \right] \\
&= E_0 \left[ (1 + \Lambda(X_1)) \left| \frac{1}{\xi_2} (e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]} - e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}) \right. \right. \\
&\quad \left. \left. - \frac{\xi_1}{\xi_2^2} (E_0[e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}] - E_0[e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}]) \right| \right] \quad (48)
\end{aligned}$$

by using mean value theorem, where  $(\xi_1, \xi_2)$  lies in the line segment between  $(e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}, E_0[e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}])$  and  $(e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}, E_0[e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}])$ . By (46), we have  $\xi_2 > 1 - \epsilon$  for some small  $\epsilon > 0$ , when  $\beta$  is small enough. Moreover,  $\xi_1 \leq e^{\beta(\Lambda(X_1) + E_0[\Lambda(X)L(X)])} \leq e^{\beta\Lambda(X_1) + \beta M}$ . We have that (48) is less than or equal to

$$\begin{aligned}
&E_0 \left[ (1 + \Lambda(X_1)) \left( \sup \left| \frac{1}{\xi_2} \right| \left| e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]} - e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]} \right| \right. \right. \\
&\quad \left. \left. + \sup \left| \frac{\xi_1}{\xi_2^2} \right| \left| E_0[e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}] - E_0[e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}] \right| \right) \right] \\
&\leq \frac{1}{1 - \epsilon} E_0 \left[ (1 + \Lambda(X_1)) \left| e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]} - e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]} \right| \right] \\
&\quad + \frac{e^{\beta M}}{(1 - \epsilon)^2} E_0 \left[ (1 + \Lambda(X_1)) e^{\beta\Lambda(X_1)} \left| E_0[e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]}] - E_0[e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]}] \right| \right] \\
&\leq C E_0 \left[ (1 + \Lambda(X_1)) e^{\beta\Lambda(X_1)} \left| e^{\beta E_0[S_h(X_1, X_2)L(X_2)|X_1]} - e^{\beta E_0[S_h(X_1, X_2)L'(X_2)|X_1]} \right| \right] \\
&\leq C \beta E_0[(1 + \Lambda(X_1)) e^{2\beta\Lambda(X_1)} |E_0[S_h(X_1, X_2)L(X_2)|X_1] - E_0[S_h(X_1, X_2)L'(X_2)|X_1]|] \\
&\quad \text{by mean value theorem again} \\
&\leq C \beta E_0[(1 + \Lambda(X_1)) e^{2\beta\Lambda(X_1)} |S_h(X_1, X_2)| |L(X_2) - L'(X_2)|] \\
&\leq C \beta E_0[(1 + \Lambda(X_1)) e^{2\beta\Lambda(X_1)} (\Lambda(X_1) + \Lambda(X_2)) |L(X_2) - L'(X_2)|] \\
&= C \beta (E_0[(1 + \Lambda(X))^2] e^{2\beta\Lambda(X)}) E_0[|L(X) - L'(X)| + E_0[(1 + \Lambda(X)) e^{2\beta\Lambda(X)}] E_0[|\Lambda(X)| |L(X) - L'(X)|]) \\
&\leq C \beta \|L - L'\|_\Lambda
\end{aligned}$$

Hence when  $\beta$  is small enough, i.e.  $\alpha$  is large enough, we have  $C\beta < 1$ , and  $\mathcal{K}$  is a strict contraction. By Banach fixed point theorem, there exists a unique fixed point  $L^*$  that satisfies  $\mathcal{K}(L^*) = L^*$ .  $\square$

**Step 2: Monotonicity of the Objective Value.** We prove that the iteration associated with  $\mathcal{K}$  is monotonic in the objective value in (40) in the following sense:

**Lemma 3.** *For sufficiently large  $\alpha$  and given any  $L^{(1)}, L^{(2)} \in \mathcal{L}(M)$ , construct the sequence  $L^{(k+1)} = \mathcal{K}(L^{(k)})$  for  $k \geq 2$  by iteratively applying  $\mathcal{K}$ . Then the value of*

$$E_0[S_h(X_1, X_2)L^{(k)}(X_1)L^{(k+1)}(X_2)] - \alpha E_0[L^{(k)} \log L^{(k)}] - \alpha E_0[L^{(k+1)} \log L^{(k+1)}]$$

*is non-decreasing in  $k$ , for  $k \geq 1$ .*

*Proof.* By Proposition 1, for any  $k \geq 2$ ,  $L^{(k+1)}$  is a maximizer of (25) in Section 4, with cost function  $E_0[S_h(X, X_2)L^{(k)}(X_2)|X_1 = x]$ . Hence, for any  $k \geq 1$ ,

$$\begin{aligned}
& E_0[S_h(X_1, X_2)L^{(k)}(X_1)L^{(k+1)}(X_2)] - \alpha E_0[L^{(k)} \log L^{(k)}] - \alpha E_0[L^{(k+1)} \log L^{(k+1)}] \\
&= E_0[E_0[S_h(X_1, X_2)L^{(k+1)}(X_2)|X_1]L^{(k)}(X_1)] - \alpha E_0[L^{(k)} \log L^{(k)}] - \alpha E_0[L^{(k+1)} \log L^{(k+1)}] \\
&\leq E_0[E_0[S_h(X_1, X_2)L^{(k+1)}(X_2)|X_1]L^{(k+2)}(X_1)] - \alpha E_0[L^{(k+2)} \log L^{(k+2)}] - \alpha E_0[L^{(k+1)} \log L^{(k+1)}] \\
&= E_0[S_h(X_1, X_2)L^{(k+1)}(X_2)L^{(k+2)}(X_1)] - \alpha E_0[L^{(k+2)} \log L^{(k+2)}] - \alpha E_0[L^{(k+1)} \log L^{(k+1)}] \\
&= E_0[S_h(X_1, X_2)L^{(k+1)}(X_1)L^{(k+2)}(X_2)] - \alpha E_0[L^{(k+1)} \log L^{(k+1)}] - \alpha E_0[L^{(k+2)} \log L^{(k+2)}]
\end{aligned}$$

by using Proposition 1 in the inequality and the symmetry of  $S_h$  in the last equality. This concludes the lemma.  $\square$

**Step 3: Convergence of the Objective Value to the Optimum.** We argue that the following convergence holds:

**Lemma 4.** *When  $\alpha$  is sufficiently large, for any  $L^{(1)}, L^{(2)} \in \mathcal{L}(M)$ , we have*

$$\begin{aligned}
& E_0[S_h(X_1, X_2)L^{(k)}(X_1)L^{(k+1)}(X_2)] - \alpha E_0[L^{(k)} \log L^{(k)}] - \alpha E_0[L^{(k+1)} \log L^{(k+1)}] \\
&\rightarrow E_0[S_h(X_1, X_2)L^*(X_1)L^*(X_2)] - 2\alpha E_0[L^* \log L^*]
\end{aligned} \tag{49}$$

where  $L^{(k)}$  is defined as in Lemma 3, and  $L^*$  is the fixed point of  $\mathcal{K}$ .

*Proof.* Let  $\beta = 1/\alpha$ . First, recognize that the fixed point theorem implies convergence of  $L^{(k)}$  to  $L^*$ , i.e.  $\|L^{(k)} - L^*\|_\Lambda \rightarrow 0$  as  $k \rightarrow \infty$ .

We consider the convergence of different parts of (49) separately. For the first term, consider

$$\begin{aligned}
& |E_0[S_h(X_1, X_2)L^{(k)}(X_1)L^{(k+1)}(X_2)] - E_0[S_h(X_1, X_2)L^*(X_1)L^*(X_2)]| \\
&\leq E_0[|S_h(X_1, X_2)| |L^{(k)}(X_1)L^{(k+1)}(X_2) - L^*(X_1)L^*(X_2)|] \\
&\leq E_0[(\Lambda(X_1) + \Lambda(X_2))(L^{(k+1)}(X_2)|L^{(k)}(X_1) - L^*(X_1)| + L^*(X_1)|L^{(k+1)}(X_2) - L^*(X_2)|)] \\
&= E_0[\Lambda(X_1)|L^{(k)}(X_1) - L^*(X_1)|] + E_0[\Lambda(X_2)L^{(k+1)}(X_2)E_0|L^{(k)}(X_1) - L^*(X_1)|] \\
&\quad + E_0[\Lambda(X_1)L^*(X_1)E_0|L^{(k+1)}(X_2) - L^*(X_2)|] + E_0[\Lambda(X_2)|L^{(k+1)}(X_2) - L^*(X_2)|] \\
&\quad \text{by the independence of } X_1 \text{ and } X_2 \text{ and the fact that } E_0[L^{(k)}] = E_0[L^{(k+1)}] = E_0[L^*] = 1 \\
&\leq C\|L^{(k)} - L^*\|_\Lambda + C\|L^{(k+1)} - L^*\|_\Lambda \\
&\rightarrow 0
\end{aligned} \tag{50}$$

as  $k \rightarrow \infty$ .

We now consider the remaining terms in (49). For  $k \geq 3$ , by noting that  $L^{(k)} = \mathcal{K}(L^{(k-1)})$ , we have

$$\begin{aligned}
& |E_0[L^{(k)} \log L^{(k)}] - E_0[L^* \log L^*]| \\
&= |(\beta E_0[S_h(X_1, X_2)L^{(k)}(X_1)L^{(k-1)}(X_2)] - \log E_0[e^{\beta E_0[S_h(X_1, X_2)L^{(k-1)}(X_2)|X_1]}]) \\
&\quad - (\beta E_0[S_h(X_1, X_2)L^*(X_1)L^*(X_2)] - \log E_0[e^{\beta E_0[S_h(X_1, X_2)L^*(X_2)|X_1]}])| \\
&\leq \beta |E_0[S_h(X_1, X_2)L^{(k)}(X_1)L^{(k-1)}(X_2)] - E_0[S_h(X_1, X_2)L^*(X_1)L^*(X_2)]| \\
&\quad + |\log E_0[e^{\beta E_0[S_h(X_1, X_2)L^{(k-1)}(X_2)|X_1]}] - \log E_0[e^{\beta E_0[S_h(X_1, X_2)L^*(X_2)|X_1]}]|
\end{aligned} \tag{51}$$

The first term in (51) converges to 0 by the same argument as (50). For the second term, we can write, by mean value theorem, that

$$\begin{aligned} & \log E_0[e^{\beta E_0[S_h(X_1, X_2)L^{(k-1)}(X_2)|X_1]}] - \log E_0[e^{\beta E_0[S_h(X_1, X_2)L^*(X_2)|X_1]}] \\ &= \frac{1}{\xi_1} (E_0[e^{\beta E_0[S_h(X_1, X_2)L^{(k-1)}(X_2)|X_1]}] - E_0[e^{\beta E_0[S_h(X_1, X_2)L^*(X_2)|X_1]}]) \end{aligned} \quad (52)$$

where  $\xi_1$  lies between  $E_0[e^{\beta E_0[S_h(X_1, X_2)L^{(k-1)}(X_2)|X_1]}]$  and  $E_0[e^{\beta E_0[S_h(X_1, X_2)L^*(X_2)|X_1]}]$ , and hence  $\xi_1 \geq E_0[e^{-\beta \Lambda(X)}]e^{\beta E_0[\Lambda(X)L(X)]} \geq 1 - \epsilon$  for some small  $\epsilon$ , when  $\beta$  is small enough. Moreover,

$$|E_0[e^{\beta E_0[S_h(X_1, X_2)L^{(k-1)}(X_2)|X_1]}] - E_0[e^{\beta E_0[S_h(X_1, X_2)L^*(X_2)|X_1]}]| \leq \beta E_0[e^{\beta \xi_2} |S_h(X_1, X_2)| |L^{(k-1)}(X_2) - L^*(X_2)|]$$

for some  $\xi_2$  lying between  $E_0[S_h(X_1, X_2)L^{(k-1)}(X_2)|X_1]$  and  $E_0[S_h(X_1, X_2)L^*(X_2)|X_1]$ , and hence  $\xi_2 \leq \Lambda(X_1) + E_0[\Lambda(X)L(X)] \leq \Lambda(X_1) + M$ . So, much like the argument in proving Lemma 2, (52) is less than or equal to

$$\begin{aligned} & \frac{\beta e^{\beta M}}{1 - \epsilon} (E_0[\Lambda(X)e^{\beta \Lambda(X)}]E_0[|L^{(k-1)}(X) - L^*(X)|] + E_0[e^{\beta \Lambda(X)}]E_0[\Lambda(X)|L^{(k-1)}(X) - L^*(X)|]) \\ & \leq C \|L^{(k-1)} - L^*\|_\Lambda \\ & \rightarrow 0 \end{aligned}$$

as  $k \rightarrow \infty$ . We conclude the lemma.  $\square$

**Final Step.** Proposition 2 is now a simple consequence of the above steps:

*Proof of Proposition 2.* Given any  $L \in \mathcal{L}(M)$ , fix  $L^{(1)} = L^{(2)} = L$  and construct the recursion  $\{L^{(k)}\}_{k \geq 1}$  as in Lemmas 3 and 4. Combining these two lemmas will conclude that

$$E_0[S_h(X_1, X_2)L(X_1)L(X_2)] - 2\alpha E_0[L \log L] \leq E_0[S_h(X_1, X_2)L^*(X_1)L^*(X_2)] - 2\alpha E_0[L^* \log L^*]$$

and hence Proposition 2.  $\square$

### 5.3 Outline of Argument of Proposition 2 for General Time Horizon $T$ .

We discuss briefly how to generalize our argument in the previous subsection to general time horizon  $T$ . There are two natural generalizations of the operator  $\mathcal{K}$  defined in the  $T = 2$  case. One is a straightforward definition in view of (41), defined on  $\mathcal{L}(M) \rightarrow \mathcal{L}(M)$ :

$$\mathcal{K}(L)(x) = \frac{e^{g^L(x)/\alpha}}{E_0[e^{g^L(X)/\alpha}]} = \frac{e^{E_0[S_h(\mathbf{X}_T)\underline{L}_{2:T}|X_1=x]/\alpha}}{E_0[e^{E_0[S_h(\mathbf{X}_T)\underline{L}_{2:T}|X_1]/\alpha}]} \quad (53)$$

where  $S_h$  is defined in (14), and  $\underline{L}_{2:T} := \prod_{t=2}^T L(X_t)$ .

Another generalization, which is more useful in the proof, is constructed as follows. Rather than using  $S_h$ , we define a uniformly symmetric function derived from  $h$  as

$$\bar{S}_h(\mathbf{x}) = \sum_{\mathbf{x} \in \mathcal{S}_T} h(\mathbf{x}) \quad (54)$$

where  $\mathcal{S}_T$  is the symmetric group of all permutations of  $\mathbf{x} = (x_1, \dots, x_T)$ . The summation in (54) has  $T!$  number of terms. By construction, the value of  $\bar{S}_h$  is invariant to any permutations of its

arguments. In the case of  $T = 2$ ,  $\bar{S}_h$  is the same as  $S_h$  defined in (14), but the difference is apparent when  $T > 2$ .

The function  $\bar{S}_h$  is only an artifact in the proof and does not appear in either the solution or practical evaluation. The main use of  $\bar{S}_h$  is to construct an operator  $\bar{\mathcal{K}} : \mathcal{L}(M)^{T-1} \rightarrow \mathcal{L}(M)^{T-1}$  as follows. Denote  $\mathbf{L} = (L_1, \dots, L_{T-1}) \in \mathcal{L}(M)^{T-1}$ . We first define the component-wise mapping  $K : \mathcal{L}(M)^{T-1} \rightarrow \mathcal{L}(M)$  given by

$$K(L_1, \dots, L_{T-1})(x) := \frac{e^{E_0[\bar{S}_h(X, X_1, \dots, X_{T-1}) \prod_{t=1}^{T-1} L_t(X_t) | X=x] / (\alpha(T-1)!)}}{E_0[e^{E_0[\bar{S}_h(X, X_1, \dots, X_{T-1}) \prod_{t=1}^{T-1} L_t(X_t) | X] / (\alpha(T-1)!)}]} \quad (55)$$

Then for a given  $\mathbf{L}$ , define

$$\begin{aligned} \tilde{L}_1 &= K(L_1, \dots, L_{T-1}) \\ \tilde{L}_2 &= K(\tilde{L}_1, L_2, \dots, L_{T-1}) \\ \tilde{L}_3 &= K(\tilde{L}_1, \tilde{L}_2, L_3, \dots, L_{T-1}) \\ &\vdots \\ \tilde{L}_{T-1} &= K(\tilde{L}_1, \dots, \tilde{L}_{T-2}, L_{T-1}) \end{aligned} \quad (56)$$

The operator  $\bar{\mathcal{K}}$  on  $\mathcal{L}(M)^{T-1}$  is defined as

$$\bar{\mathcal{K}}(\mathbf{L}) = (\tilde{L}_1, \dots, \tilde{L}_{T-1}) \quad (57)$$

It is the second operator  $\bar{\mathcal{K}}$  that turns out to satisfy a monotone property in the objective function, which is the key in proving the general case  $T > 2$ . It also holds that each component of the fixed point in  $\bar{\mathcal{K}}$  is identical and they all coincide with the fixed point in  $\mathcal{K}$  presented in (53), the latter defined on a much smaller space. The smaller space of  $\mathcal{K}$  then becomes useful for a truncation argument in random time horizon problems.

The full argument for the general time horizon  $T > 2$  will be deferred to the appendix. Following Section 5.2, below is a summary of the main steps leading to the proof. First, we need to show the contraction property and characterize the fixed point of  $\bar{\mathcal{K}}$  with an appropriate metric:

**Lemma 5.** *With Assumption 3, when  $\alpha$  is sufficiently large, the operator  $\bar{\mathcal{K}} : \mathcal{L}(M)^{T-1} \rightarrow \mathcal{L}(M)^{T-1}$  defined in (57) is well-defined, closed and a contraction, using the metric  $d(\cdot, \cdot) : \mathcal{L}(M)^{T-1} \times \mathcal{L}(M)^{T-1} \rightarrow \mathbb{R}_+$  defined as*

$$d(\mathbf{L}, \mathbf{L}') = \max_{t=1, \dots, T} E_0[(1 + \Lambda(X)) |L_t(X) - L'_t(X)|]$$

where  $\mathbf{L} = (L_1, \dots, L_{T-1}) \in \mathcal{L}(M)^{T-1}$  and  $\mathbf{L}' = (L'_1, \dots, L'_{T-1}) \in \mathcal{L}(M)^{T-1}$ . Hence there exists a unique fixed point  $\mathbf{L}^*$  that satisfies  $\mathcal{K}(\mathbf{L}^*) = \mathbf{L}^*$ . Moreover, all components of  $\mathbf{L}^*$  are identical.

The property that the fixed point  $\mathbf{L}^*$  has identical components will conclude a convergence result for the component-wise iteration defined in (55):

**Corollary 1.** *With Assumption 3 and sufficiently large  $\alpha$ , starting from any  $L^{(1)}, \dots, L^{(T-1)} \in \mathcal{L}(M)$ , the iteration  $L^{(k)} = K(L^{(k-T+1)}, \dots, L^{(k-1)})$  where  $K : \mathcal{L}(M)^{T-1} \rightarrow \mathcal{L}$  is as defined in (55), for  $k \geq T$ , converges to  $L^*$  in  $\|\cdot\|_\Lambda$ -norm, where  $L^*$  is the identical component of the fixed point  $\mathbf{L}^*$  of  $\bar{\mathcal{K}}$ .*

We shall consider the objective in (40) multiplied by  $T!$ , i.e.

$$T!(E_0[h(\mathbf{X}_T)\underline{L}_T] - \alpha E_0[L \log L]) = E_0[\bar{S}_h(\mathbf{X}_T)\underline{L}_T] - \alpha T! E_0[L \log L]. \quad (58)$$

The component-wise mapping  $K$  possesses a monotonicity property on this scaled objective:

**Lemma 6.** For sufficiently large  $\alpha$ , starting from any  $L^{(1)}, \dots, L^{(T)} \in \mathcal{L}(M)$ , construct the sequence  $L^{(k+1)} = K(L^{(k-T+2)}, \dots, L^{(k)})$  via the component-wise iteration  $K$ , for  $k \geq T$ . Then

$$E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^{(k+t-1)}(X_t) \right] - \alpha(T-1)! \sum_{t=1}^T E_0[L^{(k+t-1)} \log L^{(k+t-1)}] \quad (59)$$

is non-decreasing in  $k$ , for  $k \geq 1$ .

Finally, we show the convergence of (59) to the scaled objective in (58) evaluated at any component of the fixed point of  $\bar{K}$ :

**Lemma 7.** For sufficiently large  $\alpha$ , starting from any  $L^{(1)}, \dots, L^{(T)} \in \mathcal{L}(M)$ , we have

$$\begin{aligned} & E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^{(k+t-1)}(X_t) \right] - \alpha(T-1)! \sum_{t=1}^T E_0[L^{(k+t-1)} \log L^{(k+t-1)}] \\ \rightarrow & E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^*(X_t) \right] - \alpha T! E_0[L^* \log L^*] \end{aligned}$$

where  $L^{(k)}$  is defined by the same recursion in Lemma 6, and  $L^*$  is any (identical) component of  $\mathbf{L}^* \in \mathcal{L}(M)^{T-1}$ , the fixed point of  $\bar{K}$  defined in (57).

As in Section 5.2, given any  $L \in \mathcal{L}(M)$ , Lemmas 6 and 7 will conclude that

$$E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L(X_t) \right] - \alpha(T-1)! \sum_{t=1}^T E_0[L \log L] \leq E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^*(X_t) \right] - \alpha T! E_0[L^* \log L^*]$$

by defining  $L^{(1)} = \dots = L^{(T)} = L$  and using the recursion defined in the lemmas. Here  $L^*$  is the identical component of the fixed point of  $\bar{K}$ . Moreover,  $\bar{K}(\mathbf{L}^*) = \mathbf{L}^*$  entails setting  $L^* = K(L^*, \dots, L^*)$  for  $K$  defined in (55), which concludes that  $L^*$  satisfies (41). This concludes Proposition 2.

## 5.4 Duality Theorem and Asymptotic Expansion

Since the primal objective for the case  $T > 1$  is not necessarily concave, it is not immediate that the dual problem (39) provides the same optimal value as the primal problem (37). The following theorem, from Theorem 1 in Chapter 8, p. 220 in [39], gives a sufficient characterization over general vector space that avoids convexity. We state in the notation of this paper:

**Proposition 3** (a.k.a. Chapter 8, Theorem 1 in [39]). *With Assumptions 3 and 4 in hold, suppose there is an  $\alpha^*$ , with  $\alpha^* \geq 0$ , and an  $L^* \in \mathcal{L}(M)$  such that*

$$E_0[h(\mathbf{X}_T) \underline{L}_T^*] - \alpha^* E_0[L^* \log L^*] \geq E_0[h(\mathbf{X}_T) \underline{L}_T] - \alpha^* E_0[L \log L] \quad (60)$$

for all  $L \in \mathcal{L}(M)$ . Then  $L^*$  solves

$$\begin{aligned} & \max && E_0[h(\mathbf{X}_T) \underline{L}_T] \\ & \text{subject to} && E_0[L \log L] \leq E_0[L^* \log L^*] \\ & && L \in \mathcal{L}(M). \end{aligned}$$

For any sufficiently large  $\alpha^* > 0$ , we have already obtained an  $L^*$  (parametrized by  $\alpha^*$ ) that satisfies (60), by Proposition 2. If we can find, for each  $\eta$  in a neighborhood of 0, a large enough  $\alpha^*$  such that  $E[L^* \log L^*] = \eta$ , then this  $L^*$  will be an optimal solution to the optimization problem (37) in view of Proposition 3. The condition  $\eta = E_0[L^* \log L^*]$  is the complementary slackness condition used in the proof of Theorem 1. In fact, similar to that proof, to find an  $\alpha^*$  that satisfies  $E[L^* \log L^*] = \eta$ , we carry out an asymptotic expansion of  $E[L^* \log L^*]$  in terms of  $\alpha^*$ , and invert the expansion to express  $\alpha^*$  in terms of  $\eta$ . This will verify the optimality of  $L^*$  for (37) by Proposition 3. After that, another expansion of the objective function  $E_0[h(\mathbf{X}_T)\underline{L}_T^*]$  in terms of  $\alpha^*$  and hence  $\eta$  will then give rise to the nonparametric derivatives.

The details of the aforementioned asymptotic expansions that arrive at Theorem 2 will be left to the appendix. Below we will sketch the essential steps. The starting point is a quadratic approximation of  $L^*$ :

$$L^*(x) = 1 + \frac{1}{\alpha^*}(g(x) - E_0[g(X)]) + \frac{1}{\alpha^{*2}}V(x) + \dots$$

where we define  $V(x) := W(x) - E_0[W(X)] + \frac{1}{2}((g(x) - E_0[g(X)])^2 - E_0[(g(X) - E_0[g(X)])^2])$  and  $W(x) := \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r \neq t}} E_0[h(\mathbf{X}_T)(g(X_r) - E_0[g(X)])|X_t = x]$ . With this approximation, we will arrive at the following expansions:

**Relation between  $\alpha^*$  and  $\eta$ :** We can expand

$$\begin{aligned} \eta &= E_0[L^* \log L^*] \\ &= \frac{1}{2\alpha^{*2}}Var_0(g(X)) + \frac{1}{\alpha^{*3}} \left[ E_0[g(X)V(X)] - \frac{1}{6}\kappa_3(g(X)) \right] + O\left(\frac{1}{\alpha^{*4}}\right). \end{aligned} \quad (61)$$

Under Assumption 4, we can invert (61) to get

$$\frac{1}{\alpha^*} = \sqrt{\frac{2\eta}{Var_0(g(X))}} - \frac{2\eta(E_0[g(X)V(X)] - (1/6)\kappa_3(g(X)))}{(Var_0(g(X)))^2} + O(\eta^{3/2}). \quad (62)$$

This verifies the condition in Proposition 3, and concludes that  $L^*$ , for some large enough  $\alpha^* > 0$ , is an optimal solution to the program (37) when  $\eta$  is small enough.

**Relation between the objective value and  $\alpha^*$ :** The primal optimal objective value in (37) can be expanded as

$$E_0[h(\mathbf{X}_T)\underline{L}_T^*] = E_0[h(\mathbf{X}_T)] + \frac{1}{\alpha^*}Var_0(g(X)) + \frac{1}{\alpha^{*2}} \left[ \frac{\nu}{2} + E_0[g(X)V(X)] \right] + O\left(\frac{1}{\alpha^{*3}}\right)$$

which, combining with (62), will become

$$E_0[h(\mathbf{X}_T)] + \sqrt{2Var_0(g(X))}\eta + \frac{\eta}{Var_0(g(X))} \left[ \frac{1}{3}\kappa_3(g(X)) + \nu \right] + O(\eta^{3/2}).$$

This will conclude Theorem 2.

### 5.5 Extension to Random Time Horizon Problems

In this subsection we shall discuss the extension to random time horizon problems, under Assumption 5. This will be indicative of the argument under the alternate Assumption 6, which requires more technicality and will be left to the appendix. First, the formulation (17) can be written in terms of likelihood ratio:

$$\begin{aligned} & \max && E_0[h(\mathbf{X}_\tau)\underline{L}_\tau] \\ & \text{subject to} && E_0[L(X) \log L(X)] \leq \eta \\ & && L \in \mathcal{L} \end{aligned} \quad (63)$$

where  $\underline{L}_\tau = \prod_{t=1}^\tau L(X_t)$  comes from the martingale property of sequential likelihood ratios. We shall explain how to leverage our result in the last subsection to obtain the expressions in Theorem 3. Under Assumption 5,  $\tau \leq T$  a.s., for some  $T > 0$ . Hence the objective in (63) can also be written as  $E_0[h(\mathbf{X}_\tau)\underline{L}_\tau] = E_0[h(\mathbf{X}_\tau)\underline{L}_T]$ , again by the martingale property of  $\underline{L}_t$ .

This immediately falls back into the framework of Theorem 2, with the cost function now being  $h(\mathbf{X}_\tau)$ . For this particular cost function, we analyze  $g(x)$  and  $G(x, y)$  in Theorem 2 and argue that they are indeed in the form stated in Theorem 3. We can write

$$g(x) = \sum_{t=1}^T E_0[h(\mathbf{X}_\tau)|X_t = x] = \sum_{t=1}^T E_0[h(\mathbf{X}_\tau); \tau \geq t|X_t = x] + \sum_{t=1}^T E_0[h(\mathbf{X}_\tau); \tau < t|X_t = x] \quad (64)$$

Consider the second summation in (64). Since  $h(\mathbf{X}_\tau)I(\tau < t)$  is  $\mathcal{F}_{t-1}$ -measurable, it is independent of  $X_t$ . As a result, the second summation in (64) is constant. Similarly, we can write

$$\begin{aligned} G(x, y) &= \sum_{t=1}^T \sum_{\substack{s=1, \dots, T \\ s \neq t}} E_0[h(\mathbf{X}_\tau)|X_t = x, X_s = y] \\ &= \sum_{t=1}^T \sum_{\substack{s=1, \dots, T \\ s \neq t}} E_0[h(\mathbf{X}_\tau); \tau \geq t \wedge s|X_t = x, X_s = y] + \sum_{t=1}^T \sum_{\substack{s=1, \dots, T \\ s \neq t}} E_0[h(\mathbf{X}_\tau); \tau < t \wedge s|X_t = x, X_s = y] \end{aligned} \quad (65)$$

and the second summation in (65) is again a constant. From discussion point 3 in Section 3.2, since the first and second order nonparametric derivatives are translation invariant to  $g(x)$  and  $G(x, y)$ , Theorem 3 follows immediately.

Along the same line of observations, we can also characterize the optimal change of measure to the Lagrangian relaxation of (63) analogous to Proposition 2. For convenience, denote  $\underline{L}_s^t = \prod_{\substack{1 \leq r \leq s \\ r \neq t}} L(X_r)$ , i.e. the product of likelihood ratios, but with the  $t$ -th step identical to 1. With a cost function  $h(\mathbf{X}_\tau)$ , the function  $g^L(x)$  in Proposition 2 is now calculated to be

$$\begin{aligned} g^L(x) &= \sum_{t=1}^T g_t^L(x) = \sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_T^t|X_t = x] \\ &= \sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_T^t; \tau \geq t|X_t = x] + \sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_T^t; \tau < t|X_t = x]. \end{aligned} \quad (66)$$

For the first summation in (66), by conditioning on  $\mathcal{F}_\tau$  (or by breaking down  $I(\tau \geq t)$  into a summation in time steps  $r \geq t$  followed by conditioning each on  $\mathcal{F}_r$ ), we have

$$\sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_T^t; \tau \geq t|X_t] = \sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_\tau^t; \tau \geq t|X_t].$$

The main observation is that the second summation in (66) is independent of  $X_t$ . This is because

$$E_0[h(\mathbf{X}_\tau)\underline{L}_T^t; \tau < t|X_t] = E_0[h(\mathbf{X}_\tau)E_0[\underline{L}_T^t|\mathcal{F}_t]; \tau < t|X_t] = E_0[h(\mathbf{X}_\tau)\underline{L}_{t-1}^t; \tau < t|X_t]$$

and  $h(\mathbf{X}_\tau)\underline{L}_{t-1}^t I(\tau < t)$  is  $\mathcal{F}_{t-1}$ -measurable, hence independent of  $X_t$ . The second summation of (66) is therefore a constant equal to  $\sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_{t-1}^t; \tau < t]$ . The fixed point equation that guides the optimal change of measure is

$$\begin{aligned} L(x) &\propto e^{g^L(X)/\alpha} = \exp \left\{ \frac{1}{\alpha} \left( \sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_\tau^t; \tau \geq t|X_t = x] + \sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_{t-1}^t; \tau < t] \right) \right\} \\ &\propto \exp \left\{ \frac{1}{\alpha} \sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_\tau^t; \tau \geq t|X_t = x] \right\}. \end{aligned}$$

Comparing to Proposition 2, then,  $g^L(x)$  is replaced by  $\sum_{t=1}^T E_0[h(\mathbf{X}_\tau)\underline{L}_\tau^t; \tau \geq t|X_t = x]$  in the optimal change of measure in the context of bounded stopping time.

The case when Assumption 6 holds instead of Assumption 5 is more involved. The main idea is to consider a sequence of truncated random time  $\tau \wedge T$ ,  $T = 1, 2, \dots$ . We then set up a sequence of contraction maps  $K_T(\cdot)$  that is associated with the cost function  $h(\mathbf{X}_{\tau \wedge T})$ , and show that this sequence of maps converges to a limiting map  $K(\cdot)$ , yielding the stated result. Details will be provided in the appendix.

## 6 Bounds on Parametric Derivatives

Let us discuss how the minimization problem (2) is addressed by an easy adaptation and how the two problems (1) and (2) together give the range for all eligible parametric derivatives.

### 6.1 Minimization Formulation

All the results in Sections 3, 4 and 5 can be easily extended to the minimization formulation in (2). One merely has to replace  $h$  by  $-h$  in all the theorems. This results in a change of sign while keeping the same magnitude for the first order derivative, and keeping both the same sign and magnitude for the second order derivative, for all Theorems 1, 2 and 3. For example, the expansion in Theorem 2 becomes

$$E_{f^*}[h(\mathbf{X}_T)] = E_0[h(\mathbf{X}_T)] - \sqrt{2\text{Var}_0(g(X))}\eta^{1/2} + \frac{1}{\text{Var}_0(g(X))} \left( \frac{1}{3}\kappa_3(g(X)) + \nu \right) \eta + O(\eta^{3/2})$$

where  $g(X)$  and  $\nu$  are defined as before. In Theorem 1, moreover, the  $\beta^*$  in (5) is chosen as the unique negative instead of positive solution in the minimization formulation.

In terms of methodology, supposing one insists on using  $h$  instead of  $-h$  and carries out the minimization program directly, the Lagrangian relaxations of all the formulations in Sections 4 and 5 will become max min problems instead of min max problems. This will then lead to optimal changes of measure that involve  $\alpha < 0$ , where  $\alpha$  is the negative dual variable. Such analysis will yield the same conclusions as above.

## 6.2 Parametric Dominance

Because of the robust formulations (1) and (2), the proposed nonparametric derivatives dominate any parametric derivatives in the following sense:

**Proposition 4.** *Suppose  $P_0$  lies in a parametric family  $P^\theta$  with  $\theta \in \Theta \subset \mathbb{R}$ , say  $P_0 = P^{\theta_0}$  where  $\theta_0 \in \Theta^\circ$ . Denote  $E^\theta[\cdot]$  as the expectation under  $P^\theta$ . Assume that*

1.  $P^\theta$  is absolutely continuous with respect to  $P^{\theta_0}$  for  $\theta$  in a neighborhood of  $\theta_0$ .
2.  $D(\theta, \theta_0) := D(P^\theta \| P^{\theta_0}) \rightarrow 0$  as  $\theta \rightarrow \theta_0$ .
3. For any  $\eta$  in a neighborhood of 0,  $D(\theta, \theta_0) = \eta$  has two solutions  $\theta^+(\eta) > \theta_0$  and  $\theta^-(\eta) < \theta_0$ ; moreover,

$$\left. \frac{d}{d\theta} \sqrt{D(\theta, \theta_0)} \right|_{\theta=\theta_0^+} > 0$$

and

$$\left. \frac{d}{d\theta} \sqrt{D(\theta, \theta_0)} \right|_{\theta=\theta_0^-} < 0.$$

4.  $\left. \frac{d}{d\theta} E^\theta[h(\mathbf{X})] \right|_{\theta=\theta_0}$  exists.

Then

$$\left| \left. \frac{d}{d\theta} E^\theta[h(\mathbf{X})] \right|_{\theta=\theta_0^\pm} \right| \leq \sqrt{2\text{Var}_0(\zeta(X))}$$

where  $\zeta$  is the function  $h$ ,  $g$  or  $\tilde{g}$  in Theorems 1, 2 and 3 respectively, depending on the structure of  $\mathbf{X}$  that is stated in each theorem under the corresponding assumptions.

This proposition is an intuitive testament that our nonparametric derivative, as a result of the optimization formulation, dominates any parametric derivative that moves along the admissible directions allowed in the model space in consideration. The proof is merely a simple application of the first principle of differentiation.

*Proof.* Consider the general formulations (1) and (2), with  $P_0 = P^{\theta_0}$ . Denote  $E_{f^+(\eta)}[h(\mathbf{X})]$  as the optimal value of (1) and  $E_{f^-(\eta)}[h(\mathbf{X})]$  as the optimal value of (2), when  $\eta$  is in a neighborhood of 0. Under our assumptions,  $P^{\theta^\pm(\eta)}$  with  $D(\theta^\pm(\eta), \theta_0) = \eta$  are feasible solutions to both programs and hence the quantity  $E^{\theta^\pm(\eta)}[h(\mathbf{X})]$  satisfies  $E^{\theta^\pm(\eta)}[h(\mathbf{X})] \leq E_{f^+(\eta)}[h(\mathbf{X})]$  and  $E^{\theta^\pm(\eta)}[h(\mathbf{X})] \geq E_{f^-(\eta)}[h(\mathbf{X})]$ . This implies

$$\frac{E_{f^-(\eta)}[h(\mathbf{X})] - E_0[h(\mathbf{X})]}{\sqrt{\eta}} \leq \frac{E^{\theta^\pm(\eta)}[h(\mathbf{X})] - E_0[h(\mathbf{X})]}{\sqrt{\eta}} \leq \frac{E_{f^+(\eta)}[h(\mathbf{X})] - E_0[h(\mathbf{X})]}{\sqrt{\eta}}$$

Taking the limit as  $\sqrt{\eta} \rightarrow 0$ , the upper and lower bounds converge to our nonparametric derivatives under the corresponding assumptions in each of Theorems 1, 2 and 3. The quantities

$$\left. \frac{d}{d\sqrt{\eta}} E^{\theta^\pm(\eta)}[h(\mathbf{X})] \right|_{\sqrt{\eta}=0} = \lim_{\sqrt{\eta} \rightarrow 0} \frac{E^{\theta^\pm(\eta)}[h(\mathbf{X})] - E_0[h(\mathbf{X})]}{\sqrt{\eta}}$$

become  $\left. \frac{d}{d\theta} E^\theta[h(\mathbf{X})] \right|_{\theta=\theta_0^+}$  and  $\left. \frac{d}{d\theta} E^\theta[h(\mathbf{X})] \right|_{\theta=\theta_0^-}$  respectively, by using chain rule and implicit function theorem. This concludes the proposition.  $\square$

## 7 Numerical Implementation

In this section we demonstrate how our results, in particular Theorem 2, can be applied in practice. Let us use an example of many-server queues. Consider an FIFO queue with  $s$  number of servers, and customers arrive according to a renewal process and enact i.i.d. service times. Whenever the service capacity is full, newly arriving customers have to wait (for example, in an infinite-capacity waiting room). This generic system (and its variations) can model the dynamics of call centers, communication networks and other operation systems. In practice, the number of servers  $s$  can be big, and exact analysis is not available in general.

One common approach is to impose Markovian assumption, yet such an assumption, especially for the service time distribution, is often questionable (§III.1g in [2]). In those situations, it is useful to assess the effect on important performance measures, such as workloads and customer waiting times, when the exponential assumption is violated. For example, let us consider the mean waiting time for, say, the 100-th customer in an  $M/M/s$  queue with arrival rate 1 and service rate  $1/s$  (i.e. the system is critically loaded), starting from an empty system. The solid line in Figure 1 plots the sample average of this waiting time using 10,000 samples, for different server capacities  $s = 20, 30, 40, 50, 60$ . Also, the second and the third columns in Table 1 show these sample averages and the corresponding 95% confidence intervals.

To quantify the sensitivity of the exponential assumption for the service times, we use Theorem 2 to compute the best and the worst-case first order approximations under different levels of input model deviation  $\eta$  in terms of KL divergence. Namely, we compute the quantity  $\sqrt{2Var_0(g(X))}$  where  $P_0$  is  $Exp(1/s)$  and  $g(\cdot)$  is the symmetrization function that is computed by sequentially conditioning on the service time of customers 1 through 100, as defined in (8). Then we depict the approximation  $E_0[h(\mathbf{X}_T)] \pm \sqrt{2Var_0(g(X))}\eta$  for different levels of  $\eta$ , shown by the pairs of dotted lines in Figure 1. To get a sense,  $\eta = 0.005$  is equivalent to around 10% discrepancy in service rate if the model is *known* to lie in the family of exponential distribution; this can be seen by expressing the KL divergence in terms of service rate to see that roughly  $KL \text{ divergence} \approx (\% \text{ discrepancy in service rate})^2/2$ .

We also tabulate our point estimates of the nonparametric derivatives  $\sqrt{2Var_0(g(X))}$  and their confidence intervals in the fourth and fifth columns in Table 1. Moreover, for each  $s$ , we calculate the ratio between the nonparametric derivative estimate and the benchmark performance measure as an indicator of the relative impact of model misspecification:

$$\text{Relative model misspecification impact} := \frac{\text{Magnitude of nonparametric derivative}}{\text{Performance measure}}$$

| Number of servers | Benchmark performance measure |                | Nonparametric derivative |                  | Relative impact |
|-------------------|-------------------------------|----------------|--------------------------|------------------|-----------------|
|                   | Mean                          | C.I.           | Mean                     | C.I.             |                 |
| 20                | 5.023                         | (4.905, 5.153) | 47.840                   | (43.106, 52.574) | 9.513           |
| 30                | 3.039                         | (2.944, 3.134) | 35.437                   | (32.256, 38.618) | 11.661          |
| 40                | 1.510                         | (1.445, 1.575) | 21.504                   | (19.663, 23.345) | 14.239          |
| 50                | 0.498                         | (0.463, 0.533) | 8.255                    | (7.338, 9.171)   | 16.575          |
| 60                | 0.079                         | (0.067, 0.091) | 1.802                    | (1.490, 2.114)   | 22.702          |

Table 1: Simulation results for performance measures and first order nonparametric derivatives for the mean waiting time of the 100-th customer in  $M/M/s$  systems with different server capacities

As seen from Figure 1 and Table 1, the mean waiting time for the 100-th customer decreases




Figure 1: First order approximation of the deviation from benchmark performance measure under different input model discrepancies in terms of KL divergence

from 5.02 to 0.08, as the number of servers increases from 20 to 60. This is because the effect of critical loading reduces as more servers are available, given that the system starts from emptiness. The magnitude of the nonparametric derivative also decreases correspondingly, from 47.8 when  $s = 20$  to 1.8 when  $s = 60$ . However, the relative effect of model misspecification actually increases, from 9.51 to 22.70. This means that the larger the system capacity, the more influential it is to the mean waiting time when the exponential service time assumption is violated.

The same method can be easily adapted to test other types of performance measures and models. For example, Figure 2 and Table 2 carry out the same assessment scheme for the service time of a non-Markovian  $G/G/s$  queue with gamma arrivals and uniform service times. Here we consider a deviation from the uniform distribution for the service time. In this scenario, we see from Table 2 that both the performance measure itself and the magnitude of nonparametric derivative decrease at a slower rate compared to the  $M/M/s$  case when the number of servers increases. Namely, the mean waiting time decreases from 4.5 when  $s = 20$  to 1.9 when  $s = 60$ , while the nonparametric derivative decreases from 36.0 to 24.0. Correspondingly, the relative impact increases at a slower rate, from 8.0 to 12.7. This illustrates that the effect of model misspecification has a bigger absolute impact, but a smaller relative impact, on the waiting time in this large-scale  $G/G$  system than the Markovian counterpart. In real applications, the magnitude of  $\eta$  is chosen to capture the statistical uncertainty of the input model from past data.

Let us explain in more details about our estimation procedure. Note first that our performance measure of interest depends on both the interarrival and service times, but arrivals are auxiliary variables and hence the cost function can be written as  $E_0[h(\mathbf{X}_T, \mathbf{Y}_T)|\mathbf{X}_T]$ , where  $\mathbf{X}_T$  denotes the sequence of service times and  $\mathbf{Y}_T$  as the interarrival times (see Section 3.4). Second, note also that Assumption 3 is easily satisfied since the cost function  $E_0[h(\mathbf{X}_T, \mathbf{Y}_T)|\mathbf{X}_T] \leq \sum_{t=1}^T (E_0[Y_t] + X_t)$  and each  $E_0[Y_t] + X_t$  has exponential moment. Moreover, Assumption 4 is trivially verified by our computation that demonstrates  $g(X)$  is not a constant. Hence the assumptions in Theorem 2 is valid.

From discussion point 1 in Section 3.2, we can write  $Var_0(g(X)) = Var_0(E_0[S_h(\mathbf{X}_T, \mathbf{Y}_T)|X])$ ,




Figure 2: First order approximation of the deviation from benchmark performance measure under different input model discrepancies in terms of KL divergence

| Number of servers | Benchmark performance measure |                | Nonparametric derivative |                  | Relative impact |
|-------------------|-------------------------------|----------------|--------------------------|------------------|-----------------|
|                   | Mean                          | C.I.           | Mean                     | C.I.             |                 |
| 20                | 4.522                         | (4.427, 4.616) | 36.012                   | (33.918, 38.106) | 7.965           |
| 30                | 3.816                         | (3.725, 3.906) | 31.187                   | (29.132, 33.242) | 8.174           |
| 40                | 3.062                         | (2.981, 3.142) | 29.125                   | (27.655, 30.596) | 9.512           |
| 50                | 2.796                         | (2.718, 2.874) | 26.770                   | (24.958, 28.581) | 9.574           |
| 60                | 1.888                         | (1.820, 1.956) | 24.044                   | (21.411, 26.677) | 12.737          |

Table 2: Simulation results for performance measures and first order nonparametric derivatives for the mean waiting time of the 100-th customer in  $G/G/s$  systems with different server capacities

where  $S_h(\cdot)$  is the symmetrization sum over  $\mathbf{X}_T$  defined in (14). This quantity is therefore in the form of the variance of a conditional expectation, and we adopt an unbiased estimator from [50] that is based on analysis-of-variance (ANOVA) to estimate this quantity. This estimator takes the following form. For convenience, denote  $H := S_h(\mathbf{X}_T, \mathbf{Y}_T)$ , and we want to simulate  $Var_0(E[H|X])$ . The idea is to do a nested simulation by simulating  $X_k, k = 1, \dots, K$ , and then given each  $X_k$ , simulating  $H_{kj}, j = 1, \dots, n$ . An unbiased estimator is

$$\widehat{\sigma_M^2} = \frac{1}{K-1} \sum_{k=1}^K (\bar{H}_k - \bar{\bar{H}})^2 - \frac{1}{n} \widehat{\sigma_\epsilon^2} \quad (67)$$

where

$$\widehat{\sigma_\epsilon^2} = \frac{1}{K(n-1)} \sum_{k=1}^K \sum_{j=1}^n (H_{kj} - \bar{H}_k)^2, \quad \bar{H}_k = \frac{1}{n} \sum_{j=1}^n H_{kj} \quad \text{and} \quad \bar{\bar{H}} = \frac{1}{K} \sum_{k=1}^K \bar{H}_k$$

Next, to obtain a consistent estimate and confidence interval for  $\sqrt{2Var_0(g(X))}$ , we use the delta method and the technique of sectioning (see, for example, §III in [3]). The sampling strategy is as follows:

1. Repeat the following  $N$  times:
  - (a) Simulate  $K$  samples of  $X$ , say  $X_k = x_k, k = 1, \dots, K$ .
  - (b) For each realized  $x_k$ , simulate  $n$  samples of  $H$  given  $X = x_k$ .
  - (c) Calculate  $\widehat{\sigma}_M^2$  using (67).
2. The above procedure generates  $N$  estimators  $\widehat{\sigma}_M^2$ . Call them  $Z_l, l = 1, \dots, N$ . The final point estimator is  $\sqrt{2\bar{Z}}$ , where  $\bar{Z} = (1/N) \sum_{l=1}^N Z_l$ , and the  $1 - \alpha$  confidence interval is  $\sqrt{2} \times (\sqrt{\bar{Z}} \pm (\sigma/(2\sqrt{\bar{Z}}))t_{1-\alpha/2}/\sqrt{N})$  where  $\sigma^2 = 1/(N-1) \sum_{n=1}^N (Z_n - \bar{Z})^2$  and  $t_{1-\alpha/2}$  is the  $1 - \alpha/2$  percentile of the  $t$ -distribution with  $N - 1$  degree of freedom.

This gives a consistent estimate for  $\sqrt{2Var_0(g(X))}$  and an asymptotically valid confidence interval. In our implementation we choose  $K = 30$ ,  $n = 10$  and  $N = 20$ . The choice of the inner loop sample size  $n$  is based on a pilot estimation (see §4 in [50]).

We envision that sensitivity assessment of this sort can be applied across vast areas in engineering and management applications. As can be seen from the above example, the toolkit is applicable as long as the system can be simulated, and is methodologically identical for any input models, which provides great flexibility in applications. In inventory and revenue management for instance, it can be used to stress-test existing stocking or pricing policies and to potentially help redesign new policies that are robust to distributional uncertainty in demand and operation models. In transportation and other large-scale systems, it can be used to identify vulnerable system structures or components. Similar line of analysis can be applied to portfolio and risk management to help with investment and trading decisions. The full development of these applications shall be considered in future research.

## 8 Discussions

We close this paper by discussing a few aspects of our problem and further work.

### 8.1 Other Statistical Distances

We point out that KL divergence is not the only choice for developing nonparametric derivatives. It is however a natural choice and offers some advantages compared to other distances: It leads to the favorable exponential tilting form of the involved optimal change of measure  $L^*$ , which also arises in large deviations and rare-event problems for light-tailed systems [11]. It also has an entropy interpretation, and has been used in model calibration and derivative security valuation in the finance context (see, for example, [49]). The structurally neat form of the exponential change of measure appears to streamline the development of contraction maps and the involved convergence results shown in this paper.

Using other statistical distances will change certain aspects in the form of nonparametric derivatives and the scaling of  $\eta$  in the expansions. This arises from differences in the local behaviors of different statistical distances. As a comparison, for example,  $\chi^2$ -distance (or Euclidean distance), given by  $\chi^2(P_f, P_0) = E_0(L - 1)^2 = E_0[L^2] - 1$  will lead to an optimal change of measure satisfying the linear-type contraction map  $L(x) = 1 + (g^L(x) - E_0[g^L(X)])/(2\alpha)$  in Proposition 2. On the other hand, the empirical likelihood or the “backward” entropy [42], given by  $E_0[\log L^{-1}]$ , will give a reciprocal linear form  $L(x) = 1/(\lambda - g^L(x)/\alpha)$  where  $\lambda$  is a constant that makes  $E_0[L] = 1$ . Such changes of measure will result in different coefficients and scaling in the expansions of the involved

optimal values. Nevertheless, it appears that the “symmetrization” functions  $g(x)$  and  $G(x, y)$  and the cumulant forms come up in the nonparametric derivatives universally for many members in the Csiszar  $f$ -divergence class (§2 in [51]). This phenomenon is intuitively consistent with results from von Mises’ theory on the form of the asymptotic variance for statistical functions (§6 in [48]). We shall leave the thorough comparison of various statistical distances to future work, but shall point out that the requirement of finite exponential moment on the cost function, which by far is the most important assumption in our formulation, can potentially be removed by using other distances such as the  $\alpha$ -divergence [13, 22].

## 8.2 Simulation Output Analysis

The nonparametric derivatives developed in this paper can potentially be applied to the construction of confidence intervals subject to input model uncertainty. When the input model is estimated from data, the construction of confidence intervals for simulation outputs consists of the statistical deviations for both simulation and data collection. This is known as trace-driven or historical simulation (§2.3.2 in [40]). To use the notation in this paper, suppose one is interested in estimating  $E_0[h(\mathbf{X})]$  via simulation of size  $n$ . The exact distribution  $P_0$  for the input  $\mathbf{X}$  is unknown, but an estimated distribution, call it  $P_f$ , is inferred from collected data of size  $m$  (the form of collected data that leads to  $P_f$  can vary from case to case). Using  $P_f$ , one gets  $E_f[h(\mathbf{X})]$ , and the simulation yields  $\hat{E}_f[h(\mathbf{X})]$ . The error involved will contain

$$\hat{E}_f[h(\mathbf{X})] - E_0[h(\mathbf{X})] = (\hat{E}_f[h(\mathbf{X})] - E_f[h(\mathbf{X})]) + (E_f[h(\mathbf{X})] - E_0[h(\mathbf{X})])$$

where the first term is the error due to simulation and the second due to input uncertainty. Typically, then, the confidence interval for  $E_0[h(\mathbf{X})]$  is  $\hat{E}_f[h(\mathbf{X})] \pm z_{1-\alpha/2} \sqrt{\sigma_1^2/n + \sigma_2^2/m}$  where  $z_{1-\alpha/2}$  is the  $(1 - \alpha)$ -th percentile of normal distribution. While the simulation variance  $\sigma_1^2$  can be easily estimated, the input variance  $\sigma_2^2$  is more involved, and sectioning (§III in [3]) can be used to estimate this value in some situations.

Our nonparametric derivatives provide an alternate approach, by using the result that  $|E_f[h(\mathbf{X})] - E_0[h(\mathbf{X})]| \leq \sqrt{2\text{Var}_0(g(X))\eta} + O(\eta)$ , for any  $P_f$  that satisfies  $D(P_f||P_0) \leq \eta$ . A justified choice of  $\eta$ , which typically scales with the amount of data, can help to construct valid confidence interval by combining with the estimation of  $\text{Var}_0(g(X))$ . One possible way to do so is via empirical likelihood: since KL divergence is in the Cressie-Read family, a variant of the EL method (§3 in [42]) stipulates that, under certain regularity conditions, the associated profile likelihood is asymptotically  $\chi^2$ -distributed. This will give valid critical values for pre-specified false positive probabilities. Such development shall be left for future research.

## 8.3 Other Formulations

The formulation in this paper is the most stringent among a range of possible formulations in analyzing model misspecification. Namely, i.i.d. assumptions are held intact and uncertainty is only allowed on each identical individual distribution. On the other end of the spectrum is a formulation that allows freedom over both dependency and distributions among different random components (in a manner that is adapted to the filtration). In this case, the constraints on model misspecification in terms of KL divergence are given as  $E_0[L_t \log L_t | \mathcal{F}_{t-1}] \leq \eta$  for a sequence of likelihood ratios  $L_t$  on time steps  $t = 1, \dots, T$ . When the cost function is linear, this is equivalent to multi-stage robust control problem. General studies on the formulations in between these two extremes appear to be open.

## A Technical Proofs

### A.1 Proof of Proposition 2 for General Time Horizon $T$

We shall prove the several lemmas that lead to the proof of Proposition 2 for general time horizon  $T$  in Section 5.3. For convenience let  $C > 0$  be a generic constant.

*Proof of Lemma 5.* We prove the statement point-by-point regarding the operator  $\bar{\mathcal{K}}$ . For convenience, denote  $\bar{S}_h(X, \mathbf{X}_{T-1}) = \bar{S}_h(X, X_1, X_2, \dots, X_{T-1})$ , where  $\bar{S}_h$  is defined in (54), and  $\underline{L}_{T-1} = \prod_{t=1}^{T-1} L_t(X_t)$ . Also, denote  $\beta = 1/\alpha > 0$ , so  $\beta \rightarrow 0$  is equivalent to  $\alpha \rightarrow \infty$ .

Well-definedness and closedness: Recall the definition of  $K$  in (55), which can be written as

$$K(\mathbf{L})(x) = \frac{e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X=x]/(T-1)!}}{E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}]}$$

for any  $\mathbf{L} = (L_1, L_2, \dots, L_{T-1}) \in \mathcal{L}(M)^{T-1}$ . We shall show that, for any  $\mathbf{L} \in \mathcal{L}(M)^{T-1}$ , we have  $0 < E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}] < \infty$  and that  $K(\mathbf{L}) \in \mathcal{L}(M)$ . This will imply that, starting from any  $L_1, L_2, \dots, L_{T-1} \in \mathcal{L}(M)$ , we get a well-defined operator  $K$  and that  $\tilde{L}_1, \tilde{L}_2, \dots, \tilde{L}_{T-1}$  defined in (56) all remain in  $\mathcal{L}(M)$ . We then conclude that  $\bar{\mathcal{K}}$  is both well-defined and closed in  $\mathcal{L}(M)^{T-1}$  by the definition in (57).

Now suppose  $L_1, L_2, \dots, L_{T-1} \in \mathcal{L}(M)$ . Consider

$$\begin{aligned} E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}] &\leq E_0[e^{\beta(\Lambda(X) + \sum_{t=1}^{T-1} E_0[\Lambda(X_t)L_t(X_t)])}] \\ &= E_0[e^{\beta\Lambda(X)}]e^{\beta \sum_{t=1}^{T-1} E_0[\Lambda(X)L_t(X)]} \\ &\leq E_0[e^{\beta\Lambda(X)}]e^{\beta(T-1)M} \\ &< \infty. \end{aligned} \tag{68}$$

This also implies that  $e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!} < \infty$  a.s.. Similarly,

$$E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}] \geq E_0[e^{-\beta\Lambda(X)}]e^{-\beta(T-1)M} > 0. \tag{69}$$

Hence  $K$  is well-defined. To show closedness, consider

$$\begin{aligned} E_0[\Lambda(X)K(\mathbf{L}_{T-1})(X)] &= E_0\left[\Lambda(X)\frac{e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}}{E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}]}\right] \\ &\leq \frac{E_0[\Lambda(X)e^{\beta\Lambda(X)}]e^{2\beta(T-1)M}}{E_0[e^{-\beta\Lambda(X)}]}. \end{aligned} \tag{70}$$

Since  $E_0[\Lambda(X)e^{\beta\Lambda(X)}] \rightarrow E_0[\Lambda(X)]$  and  $E_0[e^{-\beta\Lambda(X)}] \rightarrow 1$  as  $\beta \rightarrow 0$ , (70) is bounded by  $M$  for small enough  $\beta$ , if we choose  $M > E_0[\Lambda(X)]$ . Hence  $K$  is closed in  $\mathcal{L}(M)$ .

By recursing using (56), we get that  $\bar{\mathcal{K}}$  is well-defined, and that for any  $\mathbf{L} = (L_1, \dots, L_{T-1}) \in \mathcal{L}(M)^{T-1}$ , we have  $\max_{t=1, \dots, T-1} E_0[\Lambda(X)\tilde{L}_t(X)] \leq M$ , and so  $\bar{\mathcal{K}}$  is closed in  $\mathcal{L}(M)^{T-1}$ .

Contraction: Consider, for any  $\mathbf{L} = (L_1, \dots, L_{T-1})$ ,  $\mathbf{L}' = (L'_1, \dots, L'_{T-1}) \in \mathcal{L}(M)^{T-1}$ ,

$$\begin{aligned}
& E_0[(1 + \Lambda(X))|K(\mathbf{L})(X) - K(\mathbf{L}')(X)|] \\
&= E_0 \left[ (1 + \Lambda(X)) \left| \frac{e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}}{E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}]} - \frac{e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!}}{E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!}]} \right| \right] \\
&= E_0 \left[ (1 + \Lambda(X)) \left| \frac{1}{\xi_2} (e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!} - e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!}) \right. \right. \\
&\quad \left. \left. - \frac{\xi_1}{\xi_2^2} (E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}] - E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!}]) \right| \right] \quad (71)
\end{aligned}$$

by using mean value theorem, where  $(\xi_1, \xi_2)$  lies in the line segment between  $(e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}, E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}])$  and  $(e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!}, E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!}])$ .

By (69), we have  $\xi_2 > 1 - \epsilon$  for some small  $\epsilon > 0$ , when  $\beta$  is small enough. Moreover,  $\xi_1 \leq e^{\beta(\Lambda(X)+(T-1)M)}$ . Hence, (71) is less than or equal to

$$\begin{aligned}
& E_0 \left[ (1 + \Lambda(X)) \left( \sup \left| \frac{1}{\xi_2} \right| \left| e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!} - e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!} \right| \right. \right. \\
&\quad \left. \left. + \sup \left| \frac{\xi_1}{\xi_2^2} \right| \left| E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}] - E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!}] \right| \right) \right] \\
&\leq \frac{1}{1 - \epsilon} E_0 \left[ (1 + \Lambda(X)) \left| e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!} - e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!} \right| \right] \\
&\quad + \frac{e^{\beta(T-1)M}}{(1 - \epsilon)^2} E_0 \left[ (1 + \Lambda(X)) e^{\beta \Lambda(X)} \left| E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!}] \right. \right. \\
&\quad \left. \left. - E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!}] \right| \right] \\
&\leq C E_0 \left[ (1 + \Lambda(X)) e^{\beta \Lambda(X)} \left| e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X]/(T-1)!} - e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]/(T-1)!} \right| \right] \\
&\leq C \beta E_0 \left[ (1 + \Lambda(X)) \frac{e^{2\beta \Lambda(X)}}{(T-1)!} |E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}_{T-1}|X] - E_0[\bar{S}_h(X, \mathbf{X}_{T-1})\underline{L}'_{T-1}|X]| \right] \\
&\quad \text{by mean value theorem again} \\
&\leq C \beta E_0 \left[ (1 + \Lambda(X)) \frac{e^{2\beta \Lambda(X)}}{(T-1)!} |\bar{S}_h(X, \mathbf{X}_{T-1})| |\underline{L}_{T-1} - \underline{L}'_{T-1}| \right] \\
&\leq C \beta E_0 \left[ (1 + \Lambda(X)) e^{2\beta \Lambda(X)} \left( \Lambda(X) + \sum_{t=1}^{T-1} \Lambda(X_t) \right) |\underline{L}_{T-1} - \underline{L}'_{T-1}| \right] \\
&= C \beta \left( E_0[(1 + \Lambda(X))^2 e^{2\beta \Lambda(X)}] E_0|\underline{L}_{T-1} - \underline{L}'_{T-1}| + E_0[(1 + \Lambda(X)) e^{2\beta \Lambda(X)}] \right. \\
&\quad \left. E_0 \left[ \sum_{t=1}^{T-1} \Lambda(X_t) |\underline{L}_{T-1} - \underline{L}'_{T-1}| \right] \right) \quad (72)
\end{aligned}$$

when  $\beta$  is small enough (or  $\alpha$  is large enough). Now note that

$$|\underline{L}_{T-1} - \underline{L}'_{T-1}| \leq \sum_{s=1}^{t-1} \dot{\underline{L}}_{T-1}^s |L_s(X_s) - L'_s(X_s)|$$

where each  $\dot{\underline{L}}_{T-1}^s$  is a product of either one of  $L(X_r)$  or  $L'(X_r)$  for  $r = 1, \dots, T-1, r \neq s$ . Hence (72) is less than or equal to

$$C\beta \left( E_0[(1 + \Lambda(X))^2 e^{2\beta\Lambda(X)}] \sum_{s=1}^{T-1} E_0|L_s(X) - L'_s(X)| + E_0[(1 + \Lambda(X))e^{2\beta\Lambda(X)}] \right. \\ \left. \sum_{t=1}^{T-1} \left( E_0[\Lambda(X)|L_t(X) - L'_t(X)|] + E_0[\Lambda(X)L(X)] \sum_{\substack{s=1, \dots, T-1 \\ s \neq t}} E_0|L_s(X) - L'_s(X)| \right) \right) \quad (73)$$

Now for convenience denote  $y_t = E_0[(1 + \Lambda(X))|L_t(X) - L'_t(X)|]$  and  $\tilde{y}_t = E_0[(1 + \Lambda(X))|\tilde{L}_t(X) - \tilde{L}'_t(X)|]$ , where  $\tilde{L}_t$  is defined in (56). Also denote  $\mathbf{y} = (y_1, \dots, y_{T-1})$  and  $\tilde{\mathbf{y}} = (\tilde{y}_1, \dots, \tilde{y}_{T-1})$ . Then (73) gives  $\tilde{y}_1 \leq a\mathbf{1}'\mathbf{y}$  for some  $a := a(\beta) = O(\beta)$  as  $\beta \rightarrow 0$ , and  $\mathbf{1}$  denotes the  $(T-1)$ -dimensional vector of constant 1. Then (56) implies that

$$\begin{aligned} \tilde{y}_2 &\leq a\mathbf{1}'(\tilde{y}_1, y_2, \dots, y_{T-1}) \\ \tilde{y}_3 &\leq a\mathbf{1}'(\tilde{y}_1, \tilde{y}_2, y_3, \dots, y_{T-1}) \\ &\vdots \\ \tilde{y}_{T-1} &\leq a\mathbf{1}'(\tilde{y}_1, \dots, \tilde{y}_{T-2}, y_{T-1}). \end{aligned}$$

Hence

$$\max_{t=1, \dots, T-1} \tilde{y}_t \leq \max_{t=1, \dots, T-1} \sum_{s=1}^T A_{ts} y_s$$

where  $A_{ts} := A_{ts}(\beta)$  are constants that go to 0 as  $\beta \rightarrow 0$ . Therefore, when  $\beta$  is small enough,  $d(\bar{\mathcal{K}}(\mathbf{L}_{T-1}), \bar{\mathcal{K}}(\mathbf{L}'_{T-1})) \leq wd(\mathbf{L}_{T-1}, \mathbf{L}'_{T-1})$  for some  $0 < w < 1$ , and  $\bar{\mathcal{K}}$  is a contraction. Moreover, as a consequence, starting from any initial value  $\mathbf{L}^{(1)} = \mathbf{L} \in \mathcal{L}(M)^{T-1}$ , the recursion  $\mathbf{L}^{(k+1)} = \bar{\mathcal{K}}(\mathbf{L}^{(k)})$  satisfies  $\mathbf{L}^{(k)} \xrightarrow{d} \mathbf{L}^*$  where  $\mathbf{L}^*$  is the fixed point of  $\bar{\mathcal{K}}$ .

Identical Components: It remains to show that all components of  $\mathbf{L}^*$  are the same. Denote  $\mathbf{L}^* = (L_1^*, \dots, L_{T-1}^*)$ . By definition  $\bar{\mathcal{K}}(\mathbf{L}^*) = \mathbf{L}^*$ . Using (56), we have

$$\begin{aligned} \tilde{L}_1 &= K(L_1^*, \dots, L_{T-1}^*) = L_1^* \\ \tilde{L}_2 &= K(\tilde{L}_1, L_2^*, \dots, L_{T-1}^*) = K(L_1^*, L_2^*, \dots, L_{T-1}^*) = L_2^* \\ \tilde{L}_3 &= K(\tilde{L}_1, \tilde{L}_2, L_3^*, \dots, L_{T-1}^*) = K(L_1^*, L_2^*, L_3^*, \dots, L_{T-1}^*) = L_3^* \\ &\vdots \\ \tilde{L}_{T-1} &= K(\tilde{L}_1, \dots, \tilde{L}_{T-2}, L_{T-1}^*) = K(L_1^*, \dots, L_{T-1}^*) = L_{T-1}^* \end{aligned}$$

Hence  $L_1^* = L_2^* = \dots = L_{T-1}^* = \tilde{L}_1 = \dots = \tilde{L}_{T-1} = K(L_1^*, \dots, L_{T-1}^*)$ . This concludes the lemma.  $\square$

*Proof of Corollary 1.* Note that  $\bar{\mathcal{K}}$  has a fixed point in  $\mathcal{L}(M)^{T-1}$  that has all equal components. Since convergence in  $\mathcal{L}(M)^{T-1}$  implies convergence in each component in  $\mathcal{L}(M)$  (in the  $\|\cdot\|_\Lambda$ -norm), the result follows.  $\square$

*Proof of Lemma 6.* The proof of this lemma follows the same idea as Lemma 3, with the conditioning now imposed on the variable that corresponds to the smallest index of  $L$  among the  $T-1$  candidates in consideration. More specifically, using the result in Proposition 1 and the symmetry of  $\bar{S}_h$ , we have

$$\begin{aligned}
& E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^{(k+t-1)}(X_t) \right] - \alpha(T-1)! \sum_{t=1}^T E_0[L^{(k+t-1)} \log L^{(k+t-1)}] \\
= & E_0 \left[ E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=2}^T L^{(k+t-1)}(X_t) \middle| X_1 \right] L^{(k)}(X_1) \right] - \alpha(T-1)! E_0[L^{(k)} \log L^{(k)}] \\
& - \alpha(T-1)! \sum_{t=2}^T E_0[L^{(k+t-1)} \log L^{(k+t-1)}] \\
\leq & E_0 \left[ E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=2}^T L^{(k+t-1)}(X_t) \middle| X_1 \right] L^{(k+T)}(X_1) \right] - \alpha(T-1)! E_0[L^{(k+T)} \log L^{(k+T)}] \\
& - \alpha(T-1)! \sum_{t=2}^T E_0[L^{(k+t-1)} \log L^{(k+t-1)}] \\
= & E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^{(k+t)}(X_t) \right] - \alpha(T-1)! \sum_{t=1}^T E_0[L^{(k+t)} \log L^{(k+t)}]
\end{aligned}$$

$\square$

*Proof of Lemma 7.* The proof follows closely that of Lemma 4. We consider convergence of the first and second terms of (59) separately. For the first term, consider

$$\begin{aligned}
& \left| E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^{(k+t-1)}(X_t) \right] - E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^*(X_t) \right] \right| \\
\leq & E_0 \left[ |\bar{S}_h(\mathbf{X}_T)| \left| \prod_{t=1}^T L^{(k+t-1)}(X_t) - \prod_{t=1}^T L^*(X_t) \right| \right] \\
\leq & (T-1)! E_0 \left[ \sum_{t=1}^T \Lambda(X_t) \sum_{s=1}^T \underline{\dot{L}}_T^s |L^{(k+s-1)}(X_s) - L^*(X_s)| \right] \\
& \text{where each } \underline{\dot{L}}_T^s \text{ is product of either one of } L^{(k+r-1)}(X_r) \text{ or } L^*(X_r) \text{ for } r = 1, \dots, T, r \neq s \\
\leq & C \sum_{s=1}^T E_0[(1 + \Lambda(X)) |L^{(k+s-1)}(X) - L^*(X)|] \\
\rightarrow & 0
\end{aligned} \tag{74}$$

as  $k \rightarrow \infty$ , since  $L^{(k)} \rightarrow L^*$  in  $\|\cdot\|_\Lambda$ -norm by Corollary 1.

We now consider the second term in (59). By the recursion of  $K$ , we have, for  $k \geq 1$ ,

$$\begin{aligned}
& |E_0[L^{(k+T-1)} \log L^{(k+T-1)}] - E_0[L^* \log L^*]| \\
= & \left| \left( \frac{\beta}{(T-1)!} E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^{(k+t-1)}(X_t) \right] - \log E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \prod_{t=1}^{T-1} L^{(k+t-T)}(X_t)|X]/(T-1)!}] \right) \right. \\
& \left. - \left( \frac{\beta}{(T-1)!} E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^*(X_t) \right] - \log E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \underline{L}_{T-1}^*|X]/(T-1)!}] \right) \right| \\
= & \frac{\beta}{(T-1)!} \left| E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^{(k+t-1)}(X_t) \right] - E_0 \left[ \bar{S}_h(\mathbf{X}_T) \prod_{t=1}^T L^*(X_t) \right] \right| \\
& + |\log E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \prod_{t=1}^{T-1} L^{(k+t-T)}(X_t)|X]/(T-1)!}] - \log E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \underline{L}_{T-1}^*|X]/(T-1)!}]]| \quad (75)
\end{aligned}$$

The first term in (75) converges to 0 by the same argument as in (74). For the second term, we can write, by mean value theorem, that

$$\begin{aligned}
& |\log E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \prod_{t=1}^{T-1} L^{(k+t-T)}(X_t)|X]/(T-1)!}] - \log E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \underline{L}_{T-1}^*|X]/(T-1)!}]]| \\
= & \frac{1}{\xi_1} |E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \prod_{t=1}^{T-1} L^{(k+t-1)}(X_t)|X]/(T-1)!}] - E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \underline{L}_{T-1}^*|X]/(T-1)!}]]| \quad (76)
\end{aligned}$$

where  $\xi_1$  lies between  $E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \prod_{t=1}^{T-1} L^{(k+t-1)}(X_t)|X]/(T-1)!}]$  and  $E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \underline{L}_{T-1}^*|X]/(T-1)!}]$ , and hence  $\xi_1 \geq E_0[e^{-\beta \Lambda(X)}] e^{\beta E_0[\Lambda(X) L(X)]} \geq 1 - \epsilon$  for some small  $\epsilon > 0$ , when  $\beta$  is small enough. Moreover,

$$\begin{aligned}
& |E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \prod_{t=1}^{T-1} L^{(k+t-1)}(X_t)|X]/(T-1)!}] - E_0[e^{\beta E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \underline{L}_{T-1}^*|X]/(T-1)!}]]| \\
\leq & \beta E_0 \left[ e^{\beta \xi_2} |\bar{S}_h(X, \mathbf{X}_{T-1})| \left| \prod_{t=1}^{T-1} L^{(k+t-1)}(X_t) - \underline{L}_{T-1}^* \right| \right]
\end{aligned}$$

for some  $\xi_2$  lying between  $E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \prod_{t=1}^{T-1} L^{(k+t-T)}(X_t)|X]/(T-1)!$  and  $E_0[\bar{S}_h(X, \mathbf{X}_{T-1}) \underline{L}_{T-1}^*|X]/(T-1)!$ , and hence  $\xi_2 \leq \Lambda(X) + (T-1)M$ . So, much like the argument in proving Lemma 4, (76) is less than or equal to

$$C\beta \max_{t=1, \dots, T-1} E_0[(1 + \Lambda(X)) |L^{(k+t-1)}(X) - L^*(X)|] \rightarrow 0$$

as  $k \rightarrow \infty$ . This concludes the lemma.  $\square$

## A.2 Proof of Theorem 2

We shall prove the several asymptotics discussed in Section 5.4 in order to prove Theorem 2. Denote  $\beta = 1/\alpha^* > 0$ , so  $\beta$  is small when  $\alpha^*$  is large. Then from (41) we have

$$L^*(x) = \frac{e^{\beta g^{L^*}(x)}}{E_0[e^{\beta g^{L^*}(X)}]} \quad (77)$$

where  $g^{L^*}(x) = \sum_{t=1}^T g_t^{L^*}(x) = \sum_{t=1}^T E_0[h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L^*(X_r) | X_t = x]$ . Also recall that

$$g(x) = \sum_{t=1}^T g_t(x) = \sum_{t=1}^T E_0[h(\mathbf{X}_T) | X_t = x]$$

as defined in (9). Note that  $E_0[g(X)] = TE_0[h(\mathbf{X}_T)]$ . Furthermore, let us denote, for any  $p \geq 1$ ,  $\bar{O}(\beta^p) := \bar{O}(\beta^p; x)$  as a deterministic function in  $x$  such that  $E_0[h(\mathbf{X}_T)^q \bar{O}(\beta^p; X_t)] = O(\beta^p)$  for any  $q \geq 1$  and  $t = 1, \dots, T$ , when  $\beta \rightarrow 0$ . Finally, we also let  $\psi_L(\beta) := \log E_0[e^{\beta g^L(X)}]$  for convenience.

We first give a quadratic approximation of  $L^*$  as  $\beta \rightarrow 0$  (equivalently  $\alpha^* \rightarrow \infty$ ). Then we find the relation between  $\beta$  and  $\eta$ , which verifies the optimality condition given in Proposition 3. After that we expand the objective value in terms of  $\beta$ , and hence  $\eta$ , to conclude Theorem 2.

**Asymptotic expansion of  $L^*$ :** We shall obtain a quadratic approximation of  $L^*$  by first getting a first order approximation of  $L^*$  and then iterating via the quantity  $g^{L^*}$  to get to the second order. Note that as the “logarithmic moment generating function” of  $g^{L^*}(X)$ ,

$$\begin{aligned} \psi_{L^*}(\beta) &= \log E_0[e^{\beta g^{L^*}(X)}] \\ &= \beta E_0[g^{L^*}(X)] + \frac{\beta^2}{2} \kappa_2(g^{L^*}(X)) + \frac{\beta^3}{3!} \kappa_3(g^{L^*}(X)) + O(\beta^4) \end{aligned} \quad (78)$$

where  $\kappa_2(g^{L^*}(X)) := E_0[(g^{L^*}(X) - E_0[g^{L^*}(X)])^2]$  and  $\kappa_3(g^{L^*}(X)) := [(g^{L^*}(X) - E_0[g^{L^*}(X)])^3]$ . Using (77) and (78), and the finiteness of the exponential moment of  $g^{L^*}(X)$  guaranteed by a calculation similar to (68), we have

$$\begin{aligned} L^*(x) &= \frac{e^{\beta g^{L^*}(x)}}{E_0[e^{\beta g^{L^*}(X)}]} = e^{\beta g^{L^*}(x) - \psi_{L^*}(\beta)} \\ &= 1 + \beta(g^{L^*}(x) - E_0[g^{L^*}(X)]) + \bar{O}(\beta^2) \end{aligned} \quad (79)$$

But notice that

$$\begin{aligned} g^{L^*}(x) &= \sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L^*(X_r) \middle| X_t = x \right] \\ &= \sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} (1 + \bar{O}(\beta; X_r)) \middle| X_t = x \right] \\ &= \sum_{t=1}^T E_0[h(\mathbf{X}_T) | X_t = x] + \bar{O}(\beta) \\ &= g(x) + \bar{O}(\beta) \end{aligned}$$

and hence  $E_0[g^{L^*}(X)] = E_0[g(X)] + O(\beta)$ . Consequently, from (79) we have

$$L^*(x) = 1 + \beta(g(x) - E_0[g(X)]) + \bar{O}(\beta^2) \quad (80)$$

This gives a first order approximation of  $L^*$ . Using (80), we strengthen our approximation of  $g^{L^*}$

to get

$$\begin{aligned}
g^{L^*}(x) &= \sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} (1 + \beta(g(X_r) - E_0[g(X)])) + \bar{O}(\beta^2) \right] \Big| X_t = x \\
&= g(x) + \beta \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r \neq t}} E_0[h(\mathbf{X}_T)(g(X_r) - E_0[g(X)]) | X_t = x] + O(\beta^2) \\
&= g(x) + \beta W(x) + O(\beta^2)
\end{aligned} \tag{81}$$

where we define  $W(x) := \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r \neq t}} E_0[h(\mathbf{X}_T)(g(X_r) - E_0[g(X)]) | X_t = x]$ . With (81), and using (78) again, we then strengthen the approximation of  $L^*$  to get

$$\begin{aligned}
L^*(x) &= e^{\beta g^{L^*}(x) - \psi_{L^*}(\beta)} = e^{\beta(g^{L^*}(x) - E_0[g^{L^*}(X)]) - \frac{\beta^2}{2} E_0[(g^{L^*}(X) - E_0[g^{L^*}(X)])^2] + \bar{O}(\beta^3)} \\
&= 1 + \beta(g^{L^*}(x) - E_0[g^{L^*}(X)]) + \frac{\beta^2}{2} [(g^{L^*}(x) - E_0[g^{L^*}(X)])^2 - E_0[(g^{L^*}(X) - E_0[g^{L^*}(X)])^2]] \\
&\quad + \bar{O}(\beta^3) \\
&= 1 + \beta(g(x) - E_0[g(X)]) + \beta^2 \left[ W(x) - E_0[W(X)] + \frac{1}{2} ((g(x) - E_0[g(X)])^2 - E_0[(g(X) - E_0[g(X)])^2]) \right] + \bar{O}(\beta^3) \\
&= 1 + \beta(g(x) - E_0[g(X)]) + \beta^2 V(x) + \bar{O}(\beta^3)
\end{aligned} \tag{82}$$

where we define  $V(x) := W(x) - E_0[W(X)] + \frac{1}{2} ((g(x) - E_0[g(X)])^2 - E_0[(g(X) - E_0[g(X)])^2])$ .

**Relation between  $\beta$  and  $\eta$ :** By substituting  $L^*$  depicted in (77) into  $\eta = E_0[L^* \log L^*]$ , we have

$$\eta = E_0[L^* \log L^*] = \beta E_0[g^{L^*}(X) L^*(X)] - \log E_0[e^{\beta g^{L^*}(X)}] = \beta T E_0[h(\mathbf{X}_T) \underline{L}_T^*] - \psi_{L^*}(\beta) \tag{83}$$

Using (78), we can write (83) as

$$\begin{aligned}
&\beta T E_0[h(\mathbf{X}_T) \underline{L}_T^*] - \beta E_0[g^{L^*}(X)] - \frac{\beta^2}{2} \kappa_2(g^{L^*}(X)) - \frac{\beta^3}{3!} \kappa_3(g^{L^*}(X)) + O(\beta^4) \\
&= \beta \sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L^*(X_r) (L^*(X_t) - 1) \right] - \frac{\beta^2}{2} \kappa_2(g^{L^*}(X)) - \frac{\beta^3}{3!} \kappa_3(g^{L^*}(X)) \\
&\quad + O(\beta^4)
\end{aligned} \tag{84}$$

We analyze (84) term by term. For the first term, using (82), we have

$$\begin{aligned}
& \sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} L^*(X_r)(L^*(X_t) - 1) \right] \\
&= \sum_{t=1}^T E_0 \left[ h(\mathbf{X}_T) \prod_{\substack{1 \leq r \leq T \\ r \neq t}} (1 + \beta(g(X_r) - E_0[g(X)]) + \beta^2 V(X_r) + \bar{O}(\beta^3)) \right. \\
&\quad \left. \cdot (\beta(g(X_t) - E_0[g(X)]) + \beta^2 V(X_t) + \bar{O}(\beta^3)) \right] \\
&= \beta \sum_{t=1}^T E_0[h(\mathbf{X}_T)(g(X_t) - E_0[g(X)])] + \beta^2 \left[ \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r \neq t}} E_0[h(\mathbf{X}_T)(g(X_r) - E_0[g(X)])(g(X_t) - E_0[g(X)])] \right. \\
&\quad \left. + \sum_{t=1}^T E_0[h(\mathbf{X}_T)V(X_t)] \right] + O(\beta^3) \\
&= \beta Var_0(g(X)) + \beta^2[\nu + E_0[g(X)V(X)]] + O(\beta^3) \tag{85}
\end{aligned}$$

where  $\nu$  is defined in (11). The last equality follows since

$$\begin{aligned}
& \sum_{t=1}^T E_0[h(\mathbf{X}_T)(g(X_t) - E_0[g(X)])] = \sum_{t=1}^T E_0[E_0[h(\mathbf{X}_T)|X_t](g(X_t) - E_0[g(X)])] \\
&= \sum_{t=1}^T E_0[g_t(X)(g(X) - E_0[g(X)])] = E_0[g(X)(g(X) - E_0[g(X)])] = Var_0(g(X)), \\
& \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r \neq t}} E_0[h(\mathbf{X}_T)(g(X_r) - E_0[g(X)])(g(X_t) - E_0[g(X)])] \\
&= \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r \neq t}} E_0[E_0[h(\mathbf{X}_T)|X_r, X_t](g(X_r) - E_0[g(X)])(g(X_t) - E_0[g(X)])] \\
&= E_0[G(X, Y)(g(X) - E_0[g(X)])(g(Y) - E_0[g(Y)])] = \nu
\end{aligned}$$

where  $G(X, Y)$  is defined in (12), and

$$\sum_{t=1}^T E_0[h(\mathbf{X}_T)V(X_t)] = \sum_{t=1}^T E_0[E_0[h(\mathbf{X}_T)|X_t]V(X_t)] = E_0[g(X)V(X)].$$

For the second term in (84), by using (81), we have

$$\begin{aligned}
\kappa_2(g^{L^*}(X)) &= E_0[(g^{L^*}(X) - E_0[g^{L^*}(X)])^2] \\
&= E_0[((g(X) - E_0[g(X)]) + \beta(W(X) - E_0[W(X)]) + \bar{O}(\beta^2; X))^2] \\
&= Var_0(g(X)) + 2\beta E_0[(g(X) - E_0[g(X)])(W(X) - E_0[W(X)])] + O(\beta^2) \tag{86}
\end{aligned}$$

Now notice that  $W(x)$  can be written as

$$\begin{aligned}
W(x) &= \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r \neq t}} E_0[h(\mathbf{X}_T)(g(X_r) - E_0[g(X)])|X_t = x] \\
&= \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r \neq t}} E_0[E_0[h(\mathbf{X}_T)|X_r, X_t](g(X_r) - E_0[g(X)])|X_t = x] \\
&= E_0[G(X, Y)(g(Y) - E_0[g(Y)])|X = x]
\end{aligned}$$

where  $G(X, Y)$  is defined in (12). Hence

$$\begin{aligned}
&E_0[(g(X) - E_0[g(X)])(W(X) - E_0[W(X)])] = E_0[(g(X) - E_0[g(X)])W(X)] \\
&= E_0[(g(X) - E_0[g(X)])G(X, Y)(g(Y) - E_0[g(Y)])] = \nu.
\end{aligned}$$

Consequently, (86) becomes

$$Var_0(g(X)) + 2\beta\nu + O(\beta^2). \quad (87)$$

Finally, for the third term in (84), we have

$$\kappa_3(g^{L^*}(X)) = E_0[(g(X) - E_0[g(X)])^3] + O(\beta) = \kappa_3(g(X)) + O(\beta). \quad (88)$$

Combining (85), (87) and (88), we have

$$\begin{aligned}
\eta &= \beta^2 Var_0(g(X)) + \beta^3[\nu + E_0[g(X)V(X)]] - \frac{\beta^2}{2} Var_0(g(X)) - \beta^3\nu - \frac{\beta^3}{6} \kappa_3(g(X)) + O(\beta^4) \\
&= \frac{\beta^2}{2} Var_0(g(X)) + \beta^3 \left[ E_0[g(X)V(X)] - \frac{1}{6} \kappa_3(g(X)) \right] + O(\beta^4).
\end{aligned} \quad (89)$$

Under Assumption 4, we can invert (89) to get

$$\begin{aligned}
\beta &= \sqrt{\frac{2\eta}{Var_0(g(X))}} \left( 1 + \frac{2\beta(E_0[g(X)V(X)] - (1/6)\kappa_3(g(X)))}{Var_0(g(X))} + O(\beta^2) \right)^{-1/2} \\
&= \sqrt{\frac{2\eta}{Var_0(g(X))}} - \frac{1}{2} \sqrt{\frac{2\eta}{Var_0(g(X))}} \frac{2\beta(E_0[g(X)V(X)] - (1/6)\kappa_3(g(X)))}{Var_0(g(X))} + O(\eta^{1/2}\beta^2) \\
&= \sqrt{\frac{2\eta}{Var_0(g(X))}} - \frac{2\eta(E_0[g(X)V(X)] - (1/6)\kappa_3(g(X)))}{(Var_0(g(X)))^2} + O(\eta^{3/2})
\end{aligned} \quad (90)$$

This in particular verifies the condition in Proposition 3, i.e. for any small  $\eta$ , there exists a large enough  $\alpha^* > 0$  and a corresponding  $L^*$  that satisfies (60). Consequently,  $L^*$  for some large enough  $\alpha^*$  is an optimal solution to (33).

**Relation between the objective value and  $\beta$ , and hence  $\eta$ :** Using (82) again, the optimal

objective value in (33) can be written as

$$\begin{aligned}
& E_0[h(\mathbf{X}_T)\underline{L}_T^*] \\
= & E_0 \left[ h(\mathbf{X}_T) \prod_{t=1}^T (1 + \beta(g(X_t) - E_0[g(X)]) + \beta^2 V(X_t) + \bar{O}(\beta^3; X_t)) \right] \\
= & E_0[h(\mathbf{X}_T)] + \beta \sum_{t=1}^T E_0[h(\mathbf{X}_T)(g(X_t) - E_0[g(X)])] \\
& + \beta^2 \left[ \sum_{t=1}^T \sum_{\substack{1 \leq r \leq T \\ r < t}} E_0[h(\mathbf{X}_T)(g(X_r) - E_0[g(X)])(g(X_t) - E_0[g(X)])] + \sum_{t=1}^T E_0[h(\mathbf{X}_T)V(X_t)] \right] + O(\beta^3) \\
= & E_0[h(\mathbf{X}_T)] + \beta \text{Var}_0(g(X)) + \beta^2 \left[ \frac{\nu}{2} + E_0[g(X)V(X)] \right] + O(\beta^3) \tag{91}
\end{aligned}$$

where the last equality follows from similar argument in (85). Finally, substituting (90) into (91) gives

$$\begin{aligned}
& E_0[h(\mathbf{X}_T)] + \sqrt{2\text{Var}_0(g(X))}\eta + \frac{2\eta}{\text{Var}_0(g(X))} \left[ -E_0[g(X)V(X)] + \frac{1}{6}\kappa_3(g(X)) + \frac{\nu}{2} + E_0[g(X)V(X)] \right] \\
& + O(\eta^{3/2}) \\
= & E_0[h(\mathbf{X}_T)] + \sqrt{2\text{Var}_0(g(X))}\eta + \frac{\eta}{\text{Var}_0(g(X))} \left[ \frac{1}{3}\kappa_3(g(X)) + \nu \right] + O(\eta^{3/2})
\end{aligned}$$

which coincides with Theorem 2.

### A.3 Proof of Proposition 3 under Assumption 6

We shall use a truncation argument, which involves leveraging results from finite horizon problems. We begin by observing that the operator  $\mathcal{K}$  introduced in (53) possesses similar contraction properties as the operator  $\bar{\mathcal{K}}$  in (57) in the following sense:

**Lemma 8.** *With Assumption 3 on the cost function  $h(\mathbf{X}_T)$ , for sufficiently large  $\alpha$ , the operator  $\mathcal{K} : \mathcal{L}(M) \rightarrow \mathcal{L}(M)$  defined in (53) is well-defined, closed, and a strict contraction in  $\mathcal{L}(M)$  under the metric induced by  $\|\cdot\|_\Lambda$ . Hence there exists a unique fixed point  $L^* \in \mathcal{L}(M)$  that satisfies  $\mathcal{K}(L) = L$ . Moreover,  $L^*$  is equal to each identical component of the fixed point of  $\bar{\mathcal{K}}$  defined in (57).*

*Proof.* We shall utilize our result on the operator  $\bar{\mathcal{K}}$  in Lemma 5. It is easy to check that given  $L \in \mathcal{L}(M)$ ,  $\mathcal{K}$  acted on  $L$  has the same effect as the component-wise operation  $K$ , defined in (55), acted on  $(L, \dots, L) \in \mathcal{L}(M)^{T-1}$ . In the proof of Lemma 5 we have already showed that  $K(L_1, \dots, L_{T-1})$  for any  $(L_1, \dots, L_{T-1}) \in \mathcal{L}(M)$  is well-defined, closed, and a strict contraction under  $\|\cdot\|_\Lambda$ , when  $\alpha$  is large enough (or  $\beta = 1/\alpha$  is small enough in that proof). These properties are inherited immediately to the operator  $\mathcal{K}$ .

Next, note that (41) is the fixed point equation associated with  $\mathcal{K}$ . Moreover, we have already shown in Proposition 2 that the same equation governs the fixed point of  $\bar{\mathcal{K}}$ , in the sense that the  $T-1$  components of its fixed point are all identical and satisfy (41). By the uniqueness property of fixed points, we conclude that the fixed point of  $\mathcal{K}$  coincides with each identical component of the fixed point of  $\bar{\mathcal{K}}$ .  $\square$

Now consider a cost function  $h(\mathbf{X}_\tau)$  with a random time  $\tau$  that satisfies Assumption 6. Again let  $\beta = 1/\alpha > 0$  for convenience. We introduce a sequence of truncated random time  $\tau \wedge T$ , and define  $\tilde{\mathcal{K}}_T : \mathcal{L} \rightarrow \mathcal{L}$  and  $\tilde{\mathcal{K}} : \mathcal{L} \rightarrow \mathcal{L}$  as

$$\tilde{\mathcal{K}}_T(L)(x) := \frac{e^{\beta g^{L,T}(x)}}{E_0[e^{\beta g^{L,T}(X)}]}$$

and

$$\tilde{\mathcal{K}}(L)(x) := \frac{e^{\beta g^L(x)}}{E_0[e^{\beta g^L(X)}]}$$

where

$$g^{L,T}(x) := \sum_{t=1}^T E_0[h(\mathbf{X}_{\tau \wedge T}) \underline{L}_{\tau \wedge T}^t; \tau \wedge T \geq t | X_t = x]$$

and

$$g^L(x) := \sum_{t=1}^T E_0[h(\mathbf{X}_\tau) \underline{L}_\tau^t; \tau \geq t | X_t = x].$$

Here  $\underline{L}_s^t = \prod_{r=1, \dots, s, r \neq t} L(X_r)$ . In other words,  $\tilde{\mathcal{K}}_T$  is the map identical to  $\tilde{\mathcal{K}}$  except that  $\tau$  is replaced by  $\tau \wedge T$ .

We first need the following proposition:

**Proposition 5.** *Suppose Assumption 6 is in hold. For  $\beta \leq \epsilon$  for some small  $\epsilon > 0$ , both  $\tilde{\mathcal{K}}_T$ , for any  $T \geq 1$ , and  $\tilde{\mathcal{K}}$  are well-defined, closed and a strict contraction with the same Lipschitz constant on the space  $\mathcal{L}$  equipped with the metric induced by the  $\mathcal{L}_1$ -norm  $\|L - L'\|_1 := E_0|L - L'|$ .*

*Proof.* We first consider the map  $\tilde{\mathcal{K}}$ . Recall that by Assumption 6 we have  $|h(\mathbf{X}_\tau)| \leq C$  for some constant  $C > 0$ . Consider

$$\begin{aligned} e^{\beta \sum_{t=1}^\infty E_0[h(\mathbf{X}_\tau) \underline{L}_\tau^t; \tau \geq t | X_t = x]} &\leq e^{C\beta \sum_{t=1}^\infty E_0[\underline{L}_\tau^t; \tau \geq t | X_t = x]} \\ &= e^{C\beta \sum_{t=1}^\infty E_0[\underline{L}_\tau^t; \tau \geq t]} \quad \text{since } \underline{L}_\tau^t I(\tau \geq t) \text{ is independent of } X_t \\ &= e^{C\beta \sum_{t=1}^\infty P_0(\tau \geq t)} \quad \text{since } \tau \text{ is independent of } \{X_t\}_{t \geq 1} \\ &= e^{C\beta E_0\tau} \\ &< \infty \quad \text{by Assumption 6.} \end{aligned} \tag{92}$$

Similarly,

$$e^{\beta \sum_{t=1}^\infty E_0[h(\mathbf{X}_\tau) \underline{L}_\tau^t; \tau \geq t | X_t = x]} \geq e^{-C\beta E_0\tau} > 0. \tag{93}$$

Therefore  $\tilde{\mathcal{K}}$  is well-defined and also closed in  $\mathcal{L}$ . To prove that  $\tilde{\mathcal{K}}$  is a contraction, consider, for any  $L, L' \in \mathcal{L}$ ,

$$\begin{aligned} E_0|\tilde{\mathcal{K}}(L) - \tilde{\mathcal{K}}(L')| &= E_0 \left| \frac{e^{\beta g^L(X)}}{E_0[e^{\beta g^L(X)}]} - \frac{e^{\beta g^{L'}(X)}}{E_0[e^{\beta g^{L'}(X)}]} \right| \\ &\leq E_0 \left[ \sup \left| \frac{1}{\xi_2} \right| |e^{\beta g^L(X)} - e^{\beta g^{L'}(X)}| + \sup \left| \frac{\xi_1}{\xi_2^2} \right| |E_0[e^{\beta g^L(X)}] - E_0[e^{\beta g^{L'}(X)}]| \right] \end{aligned} \tag{94}$$

by mean value theorem, where  $(\xi_1, \xi_2)$  lies in the line segment between  $(e^{\beta g^L(X)}, E_0[e^{\beta g^L(X)}])$  and  $(e^{\beta g^{L'}(X)}, E_0[e^{\beta g^{L'}(X)}])$ . By (92) and (93), we have  $\xi_1 \leq e^{C\beta E_0\tau}$  and  $\xi_2 \geq e^{-C\beta E_0\tau}$  a.s.. So (94) is less than or equal to

$$2e^{3C\beta E_0\tau} E_0 |e^{\beta g^L(X)} - e^{\beta g^{L'}(X)}| \leq 2e^{3C\beta E_0\tau} \beta E_0 |\xi| |g^L(X) - g^{L'}(X)| \quad (95)$$

by mean value theorem again, where  $\xi$  lies between  $e^{\beta g^L(X)}$  and  $e^{\beta g^{L'}(X)}$  and hence  $\xi \leq e^{C\beta E_0\tau}$  a.s.. Therefore (95) is further bounded by

$$2e^{4C\beta E_0\tau} \beta \sum_{t=1}^{\infty} E_0 [|\underline{L}_\tau^t - \underline{L}'_\tau{}^t|; \tau \geq t]. \quad (96)$$

Conditioning on  $\tau$ ,  $|\underline{L}_\tau^t - \underline{L}'_\tau{}^t| \leq \sum_{s=1, \dots, \tau, s \neq t} \dot{\underline{L}}_\tau^{t,s} |L_s - L'_s|$  where  $\dot{\underline{L}}_\tau^{t,s}$  is a product of either one  $L(X_r)$  or  $L'(X_r)$  for  $r = 1, \dots, \tau, r \neq t, s$ . Since  $\tau$  is independent of  $\{X_t\}_{t \geq 1}$  and  $\{X_t\}_{t \geq 1}$  are i.i.d., we have  $E_0[|\underline{L}_\tau^t - \underline{L}'_\tau{}^t| | \tau] \leq (\tau - 1) E_0 |L - L'|$ . Consequently, (96) is bounded by

$$2e^{4C\beta E_0\tau} \beta \sum_{t=1}^{\infty} E_0 [\tau - 1; \tau \geq t] E_0 |L - L'| = 2e^{4C\beta E_0\tau} \beta E_0 [\tau(\tau - 1)] E_0 |L - L'| \leq w E_0 |L - L'|$$

for some  $w < 1$  when  $\beta$  is small enough.

Finally, we note that the above arguments all hold with  $\tau$  replaced by  $\tau \wedge T$ , in the same range of  $\beta$  and with the same Lipschitz constant  $w$ . This concludes the proposition.  $\square$

Next we show that  $\tilde{\mathcal{K}}_T \rightarrow \tilde{\mathcal{K}}$  pointwise on  $\mathcal{L}$ :

**Proposition 6.** *Suppose Assumption 6 is in hold. For any  $L \in \mathcal{L}$ , we have  $\tilde{\mathcal{K}}_T(L) \rightarrow \tilde{\mathcal{K}}(L)$  in  $\|\cdot\|_1$ , uniformly on  $\beta \leq \epsilon$  for some small  $\epsilon > 0$ .*

*Proof.* Consider

$$\begin{aligned} E_0 |\tilde{\mathcal{K}}_T(L) - \tilde{\mathcal{K}}(L)| &= E_0 \left| \frac{e^{\beta g^{L,T}(X)}}{E_0[e^{\beta g^{L,T}(X)}]} - \frac{e^{\beta g^L(X)}}{E_0[e^{\beta g^L(X)}]} \right| \\ &\leq 2e^{4C\beta E_0\tau} \beta \sum_{t=1}^{\infty} E_0 |h(\mathbf{X}_{\tau \wedge T}) \underline{L}_{\tau \wedge T}^t I(\tau \wedge T \geq t) - h(\mathbf{X}_\tau) \underline{L}_\tau^t I(\tau \geq t)| \end{aligned} \quad (97)$$

by an argument similar to (96). Now consider

$$\begin{aligned}
& \sum_{t=1}^{\infty} E_0 |h(\mathbf{X}_{\tau \wedge T}) \underline{L}_{\tau \wedge T}^t I(\tau \wedge T \geq t) - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^t I(\tau \geq t)| \\
&= \sum_{t=1}^{\infty} E_0 [|h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^t I(\tau \geq t) - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^t I(\tau \geq t)|; \tau < T] \\
&\quad + \sum_{t=1}^{\infty} E_0 [|h(\mathbf{X}_T) \underline{L}_T^t I(T \geq t) - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^t I(\tau \geq t)|; \tau \geq T] \\
&= \sum_{t=1}^{\infty} E_0 [|h(\mathbf{X}_T) \underline{L}_T^t I(T \geq t) - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^t I(\tau \geq t)|; \tau \geq T] \\
&= \sum_{t=1}^T E_0 [|h(\mathbf{X}_T) \underline{L}_T^t - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^t|; \tau \geq T] + \sum_{t=T+1}^{\infty} E_0 [|h(\mathbf{X}_{\tau})| \underline{L}_{\tau}^t; \tau \geq t] \\
&\leq \sum_{t=1}^T E_0 [|h(\mathbf{X}_T)| |\underline{L}_T^t - \underline{L}_{\tau}^t| + |h(\mathbf{X}_T) - h(\mathbf{X}_{\tau})| \underline{L}_{\tau}^t; \tau \geq T] + C \sum_{t=T+1}^{\infty} P_0(\tau \geq t) \\
&\leq 3CTP_0(\tau \geq T) + C \sum_{t=T+1}^{\infty} P_0(\tau \geq t) \\
&\rightarrow 0
\end{aligned}$$

as  $T \rightarrow \infty$ , since  $E_0\tau < \infty$ . Hence (97) converges to 0 uniformly over  $\beta \leq \epsilon$  for some small  $\epsilon > 0$ . This concludes the proposition.  $\square$

By a simple argument on the continuity of fixed points (see, for example, Theorem 1.2 in [10]), Proposition 6 implies the following convergence result:

**Corollary 2.** *Suppose Assumption 6 is in hold. Let  $L^{(T)}$  and  $L^*$  be the fixed points of  $\tilde{\mathcal{K}}_T$  and  $\tilde{\mathcal{K}}$ . For small enough  $\beta$ , we have  $L^{(T)} \xrightarrow{\mathcal{L}_1} L^*$ .*

Finally, we show that  $L^*$  is the optimal solution to the Lagrangian relaxation of (63):

**Proposition 7.** *Under Assumption 6, the fixed point  $L^*$  of the operator  $\tilde{\mathcal{K}}$  maximizes*

$$E_0[h(\mathbf{X}_{\tau}) \underline{L}_{\tau}] - \alpha E_0[L \log L]$$

when  $\alpha$  is large enough.

Proposition 7 will initiate the same asymptotic expansion as in Section 5.4, which will give rise to the first and second order nonparametric derivatives in Theorem 3.

*Proof of Proposition 7.* We use the fact that for any fixed  $T$ ,  $L^{(T)}$  is the optimal solution to  $E_0[h(\mathbf{X}_{\tau \wedge T}) \underline{L}_{\tau \wedge T}] - \alpha E_0[L \log L]$ , as a direct consequence of Proposition 2. Hence we have the inequality

$$E_0[h(\mathbf{X}_{\tau \wedge T}) \underline{L}_{\tau \wedge T}] - \alpha E_0[L \log L] \leq E_0[h(\mathbf{X}_{\tau \wedge T}) \underline{L}_{\tau \wedge T}^{(T)}] - \alpha E_0[L^{(T)} \log L^{(T)}] \quad (98)$$

for any  $L \in \mathcal{L}$  (since  $h$  is bounded we can merely replace  $\mathcal{L}(M)$  by  $\mathcal{L}$ , i.e. putting  $M = \infty$ , in Proposition 2). Here (98) holds for any  $T \geq 1$  for  $\alpha$  uniformly large (the uniformity can be verified

using Proposition 5 and repeating the argument in the proof of Lemma 7, noting that  $\tau \wedge T \leq \tau$  a.s.). Our main argument consists of letting  $T \rightarrow \infty$  on both sides of (98).

We first show that, for any  $L \in \mathcal{L}$ , the first term on the left hand side of (98) converges to  $E_0[h(\mathbf{X}_\tau)\underline{L}_\tau]$ . Consider

$$\begin{aligned}
& |E_0[h(\mathbf{X}_{\tau \wedge T})\underline{L}_{\tau \wedge T}] - E_0[h(\mathbf{X}_\tau)\underline{L}_\tau]| \\
&= \left| \sum_{t=1}^{\infty} E_0[h(\mathbf{X}_t)\underline{L}_t]P_0(\tau \wedge T = t) - \sum_{t=1}^{\infty} E_0[h(\mathbf{X}_t)\underline{L}_t]P_0(\tau = t) \right| \\
&\leq C \sum_{t=1}^{T-1} |P_0(\tau \wedge T = t) - P_0(\tau = t)| + C|P_0(\tau \geq T) - P_0(\tau = T)| + C \sum_{t=T+1}^{\infty} |P_0(\tau \wedge T = t) - P_0(\tau = t)| \\
&= CP_0(\tau > T) + C \sum_{t=T+1}^{\infty} P_0(\tau = t) \\
&\rightarrow 0
\end{aligned}$$

as  $T \rightarrow \infty$ . Hence the left hand side of (98) converges to  $E_0[h(\mathbf{X}_\tau)\underline{L}_\tau] - \alpha E_0[L \log L]$  for any  $L \in \mathcal{L}$ . Now consider the right hand side. For the first term, consider

$$\begin{aligned}
& |E_0[h(\mathbf{X}_{\tau \wedge T})\underline{L}_{\tau \wedge T}^{(T)}] - E_0[h(\mathbf{X}_\tau)\underline{L}_\tau^*]| \\
&= \left| \sum_{t=1}^{\infty} E_0[h(\mathbf{X}_t)\underline{L}_t^{(T)}]P_0(\tau \wedge T = t) - \sum_{t=1}^{\infty} E_0[h(\mathbf{X}_t)\underline{L}_t^*]P_0(\tau = t) \right| \\
&\leq C \sum_{t=1}^{T-1} E_0|\underline{L}_t^{(T)} - \underline{L}_t^*|P_0(\tau = t) + 2C(P_0(\tau \geq T) + P(\tau = T)) + C \sum_{t=T+1}^{\infty} P_0(\tau = t) \\
&\leq C \sum_{t=1}^T tP_0(\tau = t)E_0|L^{(T)} - L^*| + 2C(P_0(\tau \geq T) + P(\tau = T)) + CP_0(\tau \geq T + 1) \\
&\quad \text{by the argument following (96)} \\
&= CE_0[\tau; \tau \leq T]E_0|L^{(T)}(X) - L^*(X)| + 2C(P_0(\tau \geq T) + P(\tau = T)) + CP_0(\tau \geq T + 1) \\
&\rightarrow 0.
\end{aligned} \tag{99}$$

Moreover, for the second term in (98), write

$$E_0[L^{(T)} \log L^{(T)}] = \beta E_0[h(\mathbf{X}_{\tau \wedge T})\underline{L}_{\tau \wedge T}^{(T)}] - \log E_0[e^{\beta g^{L^{(T)}, T}(X)}]$$

and

$$E_0[L^* \log L^*] = \beta E_0[h(\mathbf{X}_\tau)\underline{L}_\tau^*] - \log E_0[e^{\beta g^{L^*, T}(X)}]$$

by the definition of the fixed points for  $\tilde{\mathcal{K}}_T$  and  $\tilde{\mathcal{K}}$ . To prove that  $E_0[L^{(T)} \log L^{(T)}] \rightarrow E_0[L^* \log L^*]$ , we have to show that  $E_0[h(\mathbf{X}_{\tau \wedge T})\underline{L}_{\tau \wedge T}^{(T)}] \rightarrow E_0[h(\mathbf{X}_\tau)\underline{L}_\tau^*]$ , which is achieved by (99), and that  $\log E_0[e^{\beta g^{L^{(T)}, T}(X)}] \rightarrow \log E_0[e^{\beta g^{L^*, T}(X)}]$ , which we will show as follows. Consider

$$\begin{aligned}
& |\log E_0[e^{\beta g^{L^{(T)}, T}(X)}] - \log E_0[e^{\beta g^{L^*, T}(X)}]| \\
&\leq e^{C\beta E_0\tau} |E_0[e^{\beta g^{L^{(T)}, T}(X)}] - E_0[e^{\beta g^{L^*, T}(X)}]| \text{ by mean value theorem and the bound in (93)} \\
&\leq e^{2C\beta E_0\tau} \beta \sum_{t=1}^{\infty} E_0|h(\mathbf{X}_{\tau \wedge T})\underline{L}_{\tau \wedge T}^{(T)} I(\tau \wedge T \geq t) - h(\mathbf{X}_\tau)\underline{L}_\tau^* I(\tau \geq t)|
\end{aligned} \tag{100}$$

by arguments similar to (96) and (97).

Now

$$\begin{aligned}
& \sum_{t=1}^{\infty} E_0 |h(\mathbf{X}_{\tau \wedge T}) \underline{L}_{\tau \wedge T}^{(T)t} I(\tau \wedge T \geq t) - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^{*t} I(\tau \geq t)| \\
&= \sum_{t=1}^{\infty} E_0 [|h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^{(T)t} I(\tau \geq t) - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^{*t} I(\tau \geq t)|; \tau < T] \\
&\quad + \sum_{t=1}^{\infty} E_0 [|h(\mathbf{X}_T) \underline{L}_T^{(T)t} I(T \geq t) - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^{*t} I(\tau \geq t)|; \tau \geq T] \tag{101}
\end{aligned}$$

Note that the first term in (101) is bounded by

$$\begin{aligned}
C \sum_{t=1}^{\infty} E_0 [|\underline{L}_{\tau}^{(T)t} - \underline{L}_{\tau}^{*t}|; \tau \geq t, \tau < T] &\leq C \sum_{t=1}^T E_0 [\tau - 1; t \leq \tau < T] E_0 [L^{(T)} - L^*] \\
&\quad \text{by the argument following (96)} \\
&= CE_0 [\tau(\tau - 1); \tau < T] E_0 [L^{(T)} - L^*] \\
&\rightarrow 0
\end{aligned}$$

since  $E_0 \tau^2 < \infty$ . The second term in (101) can be written as

$$\begin{aligned}
& \sum_{t=1}^T E_0 [|h(\mathbf{X}_T) \underline{L}_T^{(T)t} - h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^{*t} I(\tau \geq t)|; \tau \geq T] + \sum_{t=T+1}^{\infty} E_0 [|h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^{*t}; \tau \geq t] \\
&\leq \sum_{t=1}^T E_0 [|h(\mathbf{X}_T)| |\underline{L}_T^{(T)t} - \underline{L}_{\tau}^{*t}| + |h(\mathbf{X}_T) - h(\mathbf{X}_{\tau})| \underline{L}_{\tau}^{*t}; \tau \geq T] + C \sum_{t=T+1}^{\infty} P_0(\tau \geq t) \\
&\leq 3CTP_0(\tau \geq T) + C \sum_{t=T+1}^{\infty} P_0(\tau \geq t) \\
&\rightarrow 0
\end{aligned}$$

since  $E_0 \tau < \infty$ . We therefore prove that (100) converges to 0 and the right hand side of (98) converges to  $E_0[h(\mathbf{X}_{\tau}) \underline{L}_{\tau}^*] - \alpha E_0[L^* \log L^*]$ . This concludes the proposition.  $\square$

## References

- [1] R. P. Agarwal, M. Meehan, and D. O'Regan. *Fixed Point Theory and Applications*. Cambridge University Press, 2001.
- [2] S. Asmussen. *Applied Probability and Queues*. Springer, 2nd edition, 2003.
- [3] S. Asmussen and P. W. Glynn. *Stochastic Simulation*. Springer, 2007.
- [4] C. Bandi and D. Bertsimas. Tractable stochastic analysis in high dimensions via robust optimization. *Mathematical Programming*, 134:23–70, 2012.
- [5] A. Ben-Tal, L. El Ghaoui, and A. S. Nemirovskii. *Robust Optimization*. Princeton Series in Applied Mathematics, 2009.

- [6] A. Ben-Tal and A. Nemirovski. Robust optimization: methodology and applications. *Mathematical Programming*, 92(3):453–480, 2002.
- [7] D. Bertsimas, D. Brown, and C. Caramanis. Theory and applications of robust optimization. *SIAM review*, 53(3):464–501, 2011.
- [8] J. Blanchet, P. Glynn, and H. Lam. Rare event simulation for a slotted time  $M/G/s$  model. *Queueing Systems*, 63:33–57, 2009.
- [9] J. Blanchet and C. Li. Efficient simulation for the maximum of infinite horizon discrete-time Gaussian processes. *J. Appl. Probab.*, 48:467–489, 2011.
- [10] F. F. Bonsall. *Lectures on Some Fixed Point Theorems of Functional Analysis*. Tata Institute of Fundamental Research, Bombay, India, 1962.
- [11] J. Bucklew. *Introduction to Rare Event Simulation*. Springer-Verlag, 2004.
- [12] D. R. Cox and D. V. Hinkley. *Theoretical Statistics*. Chapman & Hall/CRC, 1974.
- [13] S. Dey and S. Juneja. Incorporating fat tails in financial models using entropic divergence measures. *Working paper, Available at arXiv: <http://arxiv.org/abs/1203.0643>*, 2012.
- [14] R. Durrett. *Probability Theory and Examples*. Cambridge Series in Statistical and Probabilistic Mathematics, 4th edition, 2010.
- [15] B. L. Fox and P. W. Glynn. Replication schemes for limiting expectations. *Probability in the Engineering and Informational Sciences*, 3(03):299–318, 1989.
- [16] M. Fu. Gradient estimation. In *Chapter 19, in Handbook on Operations Research and Management Science: Simulation, S.G. Henderson and B.L. Nelson, editors*, pages 575–616. Elsevier, 2006.
- [17] M. C. Fu and J. Q. Hu. *Conditional Monte Carlo: Gradient Estimation and Optimization Applications*. Kluwer Academic Publishers, Norwell, MA, 1997.
- [18] P. Glasserman. *Gradient Estimation via Perturbation Analysis*. Kluwer Academic Publishers, Norwell, MA, 1991.
- [19] P. Glasserman. Structural conditions for perturbation analysis derivative estimation: Finite-time performance indices. *Operations Research*, 39(5):724–738, 1991.
- [20] P. Glasserman. *Monte Carlo Methods in Financial Engineering*. Springer-Verlag, New York, 2004.
- [21] P. Glasserman and X. Xu. Robust portfolio control with stochastic factor dynamics. *Working paper, Available at SSRN: <http://ssrn.com/abstract=1960723>*, 2012.
- [22] P. Glasserman and X. Xu. Robust risk measurement and model risk. *Working paper, Available at SSRN: <http://ssrn.com/abstract=2167765>*, 2012.
- [23] P. W. Glynn. Likelihood ratio gradient estimation for stochastic systems. *Commun. ACM*, 33(10):75–84, Oct. 1990.

- [24] W.-B. Gong and Y.-C. Ho. Smoothed (conditional) perturbation analysis of discrete event dynamical systems. *IEEE Transactions on Automatic Control*, 32(10):858 – 866, 1987.
- [25] F. Hampel. The influence curve and its role in robust estimation. *Journal of the American Statistical Association*, 69(346):383–393, 1974.
- [26] L. P. Hansen and T. J. Sargent. Robust control and model uncertainty. *The American Economic Review*, 91(2):pp. 60–66, 2001.
- [27] L. P. Hansen and T. J. Sargent. *Robustness*. Princeton University Press, 2008.
- [28] Y. C. Ho and X. R. Cao. Perturbation analysis and optimization of queueing networks. *Journal of Optimization Theory and Applications*, 40:559–582, 1983.
- [29] L. J. Hong. Estimating quantile sensitivities. *Operations Research*, 57(1):118–130, 2009.
- [30] A. Jain, A. Lim, and J. Shanthikumar. On the optimality of threshold control in queues with model uncertainty. *Queueing Systems*, 65:157–174, 2010.
- [31] S. Kullback and R. A. Leibler. On information and sufficiency. *The Annals of Mathematical Statistics*, 22(1):pp. 79–86, 1951.
- [32] P. L’Ecuyer. A unified view of the IPA, SF, and LR gradient estimation techniques. *Management Science*, 36(11):1364–1383, 1990.
- [33] P. L’Ecuyer. An overview of derivative estimation. In *Proceedings of the 23rd conference on Winter simulation*, WSC ’91, pages 207–217, Washington, DC, USA, 1991. IEEE Computer Society.
- [34] P. L’Ecuyer and G. Perron. On the convergence rates of IPA and FDC derivative estimators. *Operations Research*, 42(4):643–656, 1994.
- [35] A. E. B. Lim, G. J. Shanthikumar, and M. Z. J. Shen. Model uncertainty, robust optimization and learning. *Tutorials in Operations Research*, 2006.
- [36] A. E. B. Lim and J. G. Shanthikumar. Relative entropy, exponential utility, and robust dynamic pricing. *Operations Research*, 55(2):198–214, 2007.
- [37] A. E. B. Lim, J. G. Shanthikumar, and T. Watwai. Robust asset allocation with benchmarked objectives. *Mathematical Finance*, 21(4):643–679, 2011.
- [38] G. Liu and L. J. Hong. Kernel estimation of the Greeks for options with discontinuous payoffs. *Operations Research*, 59(1):96–108, 2011.
- [39] D. G. Luenberger. *Optimization by Vector Space Methods*. Wiley, 1969.
- [40] A. J. McNeil, R. Frey, and P. Embrechts. *Quantitative Risk Management: Concepts, Techniques and Tools*. Princeton Series in Finance, 2005.
- [41] M. K. Nakayama. Asymptotics of likelihood ratio derivative estimators in simulations of highly reliable Markovian systems. *Management Science*, 41(3):524–554, 1995.
- [42] A. B. Owen. *Empirical Likelihood*. Chapman & Hall/CRC, 2001.

- [43] I. Petersen, M. James, and P. Dupuis. Minimax optimal control of stochastic uncertain systems with relative entropy constraints. *IEEE Transactions on Automatic Control*, 45(3):398–412, 2000.
- [44] M. I. Reiman and A. Weiss. Sensitivity analysis for simulations via likelihood ratios. *Operations Research*, 37(5):830–844, 1989.
- [45] R. Y. Rubinstein. Sensitivity analysis and performance extrapolation for computer simulation models. *Operations Research*, 37(1):72–81, 1989.
- [46] R. Y. Rubinstein. Sensitivity analysis of discrete event systems by the push out method. *Annals of Operations Research*, 39:229–250, 1992.
- [47] R. Y. Rubinstein and A. Shapiro. *Discrete Event Systems: Sensitivity Analysis and Stochastic Optimization by the Score Function Method*. Wiley, New York, 1993.
- [48] R. J. Serfling. *Approximation Theorems of Mathematical Statistics*. Wiley, 1980.
- [49] M. Stutzer. A simple nonparametric approach to derivative security valuation. *The Journal of Finance*, 51(5):1633–1652, 1996.
- [50] Y. Sun, D. W. Apley, and J. Staum. Efficient nested simulation for estimating the variance of a conditional expectation. *Operations Research*, 59(4):998–1007, 2011.
- [51] A. B. Tsybakov. *Introduction to Nonparametric Estimation*. Springer, 2009.
- [52] Y. Wang, M. C. Fu, and S. I. Marcus. A new stochastic derivative estimator for discontinuous payoff functions with application to financial derivatives. *Operations Research*, 60(2):447–460, 2012.
- [53] M. Zazanis and R. Suri. Convergence rates of finite-difference sensitivity estimates for stochastic systems. *Operations research*, 41(4):694–703, 1993.